



Form PTO-1500 (Rev. 11-2000)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER 9013.43
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371				U.S. APPLICATION NO. (If known, see 37 CFR 1.5) 10/069082
INTERNATIONAL APPLICATION NO. PCT/GB00/03159		INTERNATIONAL FILING DATE 16 August 2000		PRIORITY DATE CLAIMED 18 August 1999
TITLE OF INVENTION Retrovirus Assay				
APPLICANT(S) FOR DO/EO/US Daniel GALBRAITH; Helena KELLY; Kenneth SMITH				
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:				
<ol style="list-style-type: none">1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.3. <input type="checkbox"/> This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below.4. <input type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (Article 31).5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2))<ol style="list-style-type: none">a. <input checked="" type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau).b. <input type="checkbox"/> has been communicated by the International Bureau.c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US).6. <input type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).<ol style="list-style-type: none">a. <input type="checkbox"/> is attached hereto.b. <input type="checkbox"/> has been previously submitted under 35 U.S.C. 154(d)(4)7. <input type="checkbox"/> Amendments to the claims of the International Application Under PCT Article 19 (35 U.S.C. 371(c)(3))<ol style="list-style-type: none">a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau).b. <input type="checkbox"/> have been communicated by the International Bureau.c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired.d. <input type="checkbox"/> have not been made and will not be made.8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).9. <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).10. <input type="checkbox"/> An English language translation of the annexes of the International Preliminary Examination Report Under PCT Article 36 (35 U.S.C. 371(c)(5)).				
Items 11 to 20 below concern document(s) or information included:				
<ol style="list-style-type: none">11. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98.12. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.13. <input checked="" type="checkbox"/> A FIRST preliminary amendment.14. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment.15. <input type="checkbox"/> A substitute specification.16. <input type="checkbox"/> A change of power of attorney and/or address letter.17. <input type="checkbox"/> A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.18. <input type="checkbox"/> A second copy of the published international application under 35 U.S.C. 154(d)(4).19. <input type="checkbox"/> A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4)20. <input type="checkbox"/> Other items or information: Written Opinion; International Search Report				

U.S. APPLICATION NO. 10/069082		INTERNATIONAL APPLICATION NO. PCT/GB00/03159	ATTORNEY DOCKET NO. 9013.43
21. <input type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492(a) (1) - (5)):			CALCULATIONS PTO USE ONLY
Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO.....			\$1040.00
International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO.....			\$890.00
International preliminary examination fee 37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO.....			\$740.00
International preliminary examination fee (37 CFR 1.482) paid to USPTO by all claims did not satisfy provisions of PCT Article 33(1)-(4).....			\$710.00
International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4).....			\$100.00
ENTER APPROPRIATE BASE FEE AMOUNT =			\$1040.00
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(c)).			\$
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	19 - 20 =	0	x \$18.00
Independent Claims	1 - 3 =	0	x \$84.00
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$280.00
TOTAL OF ABOVE CALCULATIONS =			\$1320.00
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2			\$
SUBTOTAL =			\$1320.00
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).			\$
TOTAL NATIONAL FEE =			\$1320.00
Fee for Recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +			\$
TOTAL FEES ENCLOSED =			\$1320.00
			Amount to be refunded: \$
			charged: \$
a. <input checked="" type="checkbox"/> A check in the amount of \$1320.00 to cover the above fees is enclosed.			
b. <input type="checkbox"/> Please charge my Deposit Account No. 50-0220 in the amount of \$_____ to cover the above fees. A duplicate copy of this sheet is enclosed.			
c. <input type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 50-0220. A duplicate copy of this sheet is enclosed.			
d. <input type="checkbox"/> Fees are to be charged to a credit card. WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.			
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.			
SEND ALL CORRESPONDENCE TO:			
 Kenneth D. Sibley, Reg. #31,663			
Date: February 15, 2002			
CERTIFICATE OF EXPRESS MAILING			
Express Mail Label No. EV015808902US			
Date of Deposit: February 15, 2002			
I hereby certify that this correspondence is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to: BOX PCT, Attn: DO/EO/US, Commissioner for Patents, Washington, DC 20231.			
 Vickie Diane Prior			



20792

PATENT TRADEMARK OFFICE



Rec'd PCT/PTO 30 OCT 2002

Attorney Docket No. 9013.43

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re: Galbraith, et al.
U.S. Serial No.: 10/069,082
Int'l Application No.: PCT/GB00/03159
Int'l Filing Date: August 26, 2000
For: RETROVIRUS ASSAY

October 25, 2002


Commissioner for Patents
Washington, DC 20231

STATEMENT IN SUPPORT OF FILING A
SEQUENCE LISTING UNDER 37 CFR § 1.821(f)

Sir:

Submitted herewith is a copy of the Sequence Listing in computer readable form. I hereby state that the content of the paper and computer readable copies of the Sequence listing are the same. I also hereby state as required by 37 CFR § 1.821(h) that the computer readable copy submitted concurrently herewith contains no new matter, nor does it go beyond the disclosure of the application as filed.

Respectfully submitted,


E. Michael Sajovec
Registration No. 31,793



20792

PATENT TRADEMARK OFFICE

Certificate of Mailing under 37 CFR 1.8

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Commissioner for Patents, Washington, DC 20231, on October 25, 2002.


Vickie Diane Prior

Date of Signature: October 25, 2002

IN THE UNITED STATES DESIGNATED OFFICE (DO/US)

In re: Galbraith, et al.
Serial No. To Be Assigned
Filed: Concurrently Herewith
For: RETROVIRUS ASSAY

February 15, 2002

Box PCT
Attn: DO/US
Commissioner for Patents
Washington, DC 20231

PRELIMINARY AMENDMENT

Sir:

Prior to the examination of the above application, please amend the above-identified application as follows. If any extension of time for the accompanying response or submission is required, Applicant requests that this be considered a petition therefor. The Commissioner is hereby authorized to charge any additional fee, which may be required, or credit any refund, to our Deposit Account No. 50-0220.

In the Specification:

-- RELATED APPLICATIONS

The present application claims priority from International Application No. PCT/GB00/03159, filed on 16 August 2000, which in turn claims priority from British application 9919604.0, filed on 18 August 1999, the disclosures of which are hereby incorporated herein by reference in their entirety.--

In the Claims:

Please enter the following amended claims:

13 (amended). An antiserum specific to a polypeptide fragment in accordance with Claim 1.

In re: Galbraith, et al.
International Appl. No. PCT/GB00/03159
International Filing Date: 16 August 2000
Page 2

14 (amended). A PoERV specific antibody or fragment thereof raised against a polypeptide fragment in accordance with Claim 1.

16 (amended). Use of a polypeptide fragment according to Claim 1 in detection of PoERV antibodies in a sample.

18 (amended). An assay kit for use in detection of PoERV antibodies in a sample, the kit comprising a polypeptide fragment in according with Claim 1.


19 (amended). Use of an antibody according to Claim 14, or a polypeptide fragment according to Claim 1 in therapy or diagnosis.

In re: Galbraith, et al.
International Appl. No. PCT/GB00/03159
International Filing Date: 16 August 2000
Page 3

REMARKS

Claims 1-19 are presented for examination. The above claims have been amended to better conform to U.S. practice. Applicants respectfully request substantive examination on the merits. A copy of the claims showing the changes made is attached hereto as a "Version with Markings to Show Changes Made".

Respectfully submitted,


Kenneth D. Sibley
Registration No. 31,665




20792

PATENT TRADEMARK OFFICE

CERTIFICATE OF EXPRESS MAILING

"Express Mail" mailing label number EL015808902US
Date of Deposit: February 15, 2002

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to Box PCT, Commissioner for Patents, Washington, DC 20231.


Vickie Diane Prior
Date of Signature: February 15, 2002

In re: Galbraith, et al.
International Appl. No. PCT/GB00/03159
International Filing Date: 16 August 2000
Page 4

VERSION WITH MARKINGS TO SHOW CHANGES MADE

13 (amended). An antiserum specific to a polypeptide fragment in accordance with **[any preceding claim]** Claim 1.

14 (amended). A PoERV specific antibody or fragment thereof raised against a polypeptide fragment in accordance with **[any of]** Claim[s] 1 **[to 12]**.

16 (amended). Use of a polypeptide fragment according to **[any one of]** Claim[s] 1 **[to 12]** in detection of PoERV antibodies in a sample.

18 (amended). An assay kit for use in detection of PoERV antibodies in a sample, the kit comprising a polypeptide fragment in according with **[any of]** Claim[s] 1 **[to 12]**.

19 (amended). Use of an antibody according to Claim 14, or a polypeptide fragment according to **[any one of]** Claim[s] 1 **[to 12]** in therapy or diagnosis.

infection of human cells. Since PoERV is expressed in pigs there is the potential for virus to be present in material prepared from pigs. Furthermore, as a consequence of xenotransplantation using porcine donor organs, there is
5 the possibility that the endogenous virus will be expressed in vivo and be a potential risk of PoERV infection of the patient and the general population thereafter.

A number of different types of PoERV are known, based on their genetic makeup. Types designated PERV A, PERV B,
10 and PoEV1 are described in International Patent Application WO97/40167, while types designated PERV MSL and Tsukuba are described in International Patent Application WO97/21836.

PoERV viruses comprise three genes: *gag*, *pol*, and *env*, generating GAG, POL and ENV polypeptides. It has been
15 observed that the *gag* region of the genome appears to be substantially conserved among different viruses, as well as between PoERV virus types, while the *env* region contains both conserved and non-conserved regions, which non-conserved regions are observed to vary between viral types.

It is among the objects of the present invention to provide means whereby patients and/or samples may be monitored for viral infection. It is further among the
20 objects of the present invention to provide means whereby the viral type may be determined.

According to one aspect of the present invention there is provided a PoERV polypeptide fragment, wherein said polypeptide fragment has PoERV specific antigenic or
25 immunogenic activity. Antigenic or immunogenic activity is

to be understood as capable of eliciting a PoERV specific immune response when introduced into a normal mammalian host. For example, PoERV specific antibodies are produced as a consequence.

5 According to a further aspect of the present invention there is provided an antiserum specific to a PoERV polypeptide fragment as described above.

 According to one aspect of the present invention, there is provided a fragment of a PoERV GAG polypeptide, 10 wherein said fragment has PoERV specific antigenic or immunogenic activity. Such a fragment will be referred to hereinafter as a GAG peptide, it being understood that this is distinct from native GAG protein, and may comprise only a fragment thereof, provided the GAG peptide has antigenic 15 activity. The consensus PoERV GAG polypeptide sequence is shown in Figure 1; in preferred embodiments of the invention, the GAG peptide may be selected from within this sequence.

 According to a further aspect of the present 20 invention, there is provided a fragment of a PoERV ENV polypeptide, wherein said fragment has PoERV specific antigenic or immunogenic activity. Such a fragment will be referred to hereinafter as an ENV peptide, it being understood that this is distinct from native ENV protein, 25 and may comprise only a fragment thereof, provided the ENV peptide has antigenic activity. In one embodiment, the ENV peptide may be selected from within a conserved region of the various PoERV sequences, as illustrated in Figures 2

and 3. Conserved regions are those which comprise identical and/or highly conserved amino acid sequences in different PoERV virus types; wholly conserved amino acids are indicated in Figures 2 and 3 by an asterisk beneath the amino acid, with highly conserved amino acids being indicated by a colon. Alternatively, the ENV peptide may be selected from within the non-conserved regions of the various PoERV sequences of Figures 2 and 3, in which case the ENV peptide will be specific for a particular type of PoERV. Specific examples of non-conserved type-specific ENV peptides are shown as peptides D-H and J in Figure 3. In a third embodiment, the ENV peptide may comprise both a conserved and a non-conserved region of the PoERV ENV protein, from either adjacent or non-adjacent regions of the ENV protein. Such peptides may be considered useful in simultaneous detection of any PoERV virus and a specific viral type.

In a further aspect of the present invention, there is provided a fusion GAG/ENV peptide, which peptide comprises both GAG peptide sequences and ENV peptide sequences. Such peptide may be considered useful in simultaneous detection of any PoERV virus, by means of the GAG peptide, and a specific viral type, by means of the ENV peptide.

According to a yet further aspect of the present invention there is provided antibodies specific to either GAG or ENV peptides. The antibodies may be polyclonal or monoclonal. Such antibodies can include, but are not limited to, polyclonal antibodies, monoclonal antibodies

(mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the
5 above. The Ig tails of such antibodies can be modified to reduce complement activation and Fc binding, (See, for example, European Patent No. 239400 B1, Aug. 3, 1994).

For the production of antibodies to a peptide, various host animals can be immunized by injection with a peptide,
10 or a portion thereof. Such host animals can include but are not limited to rabbits, mice, and rats, to name but a few. Various adjuvants can be used to increase the immunological response, depending on the host species, including but not limited to Freund's (complete and
15 incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guérin) and
20 *Corynebacterium parvum*.

Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of animals immunized with an antigen, such as target gene product, or an antigenic functional derivative thereof. For the
25 production of polyclonal antibodies, host animals such as those described above, can be immunized by injection with a gene product supplemented with adjuvants as also described above.

Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, can be obtained by any technique which provides for the production of antibody molecules by continuous cell lines in culture.

5 These include, but are not limited to the hybridoma technique of Kohler and Milsrein, (1975, Nature 256:495-497; and US Pat. No. 4,376,110), the human B-cell hybridoma technique (Kosbor et al., 1983, Immunology Today 4:72; Cole et al., 1983, Proc. Natl. Acad. Sci. U.S.A. 80:2026-2030),

10 and the EBV-hybridoma technique (Cole et al., 1985, Monoclonal Anti-bodies And Cancer Therapy, Alan R. Liss, Inc., pp.77-96).

Such antibodies can be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof.

15 The hybridoma producing the mAb of this invention can be cultivated in vitro or in vivo. Production of high titers of mAbs in vivo makes this the presently preferred method of production.

In addition, techniques for the production of "chimeric antibodies" (Morrison et al., 1984, Proc. Natl. Acad. Sci., 81:6851-6855; Neuberger et al., 1984, Nature, 312:604-608; Takeda et al., 1985, Nature, 314:452-454; U.S. Pat. No. 4,816,567) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with

25 genes from a human antibody molecule of appropriate biological activity can be used. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable

region derived from a murine mAb and a human immunoglobulin constant region.

Alternatively, techniques described for the production of single chain antibodies (U.S. Pat. No. 4,946,778: Bird, 1988, Science 242:423-426; Huston et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85:5879-5883; and Ward et al., 1989, Nature 334:544-546) and for making humanized monoclonal antibodies (U.S. Pat. No. 5,225,539) can be utilized to produce anti-differentially expressed or anti-pathway gene product antibodies.

Antibody fragments which recognize specific epitopes can be generated by known techniques. For example, such fragments include but are not limited to: the $F(ab')_2$ fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the $F(ab')_2$ fragments. Alternatively, Fab expression libraries can be constructed (Huse et al., 1989, Science, 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

Further aspects of the present invention provide methods of screening serum or tissue from humans or animal recipients of porcine tissue for exposure to PoERV. These methods include: use of antibodies to GAG or ENV peptides in the capture and/or detection of PoERV antigens; use of antibodies to GAG or ENV peptides in the detection of PoERV gene expression in virus infected cells by indirect immunofluorescence staining; the use of antibodies to GAG

or ENV peptides in the visualisation of PoERV virions in a sample by immuno-electron microscopy; use of GAG or ENV peptides in Western blotting for the detection of PoERV antibodies in samples from recipients of porcine-derived materials; and the use of GAG or ENV peptides in an enzyme-linked immunosorbent assay (ELISA) for the detection of PoERV antibodies.

Each of these methods may non-specifically detect any PoERV type (if GAG peptides, conserved ENV peptides or antibodies are used) or specific PoERV types, if type-specific non-conserved ENV peptides or antibodies are used.

The present invention also encompasses assay kits including GAG or ENV peptides or antibodies to such peptides, for use in the abovementioned assays. In preferred embodiments, the kits may further comprise any or all necessary preparative reagents, washing reagents, detection reagents and signal producing reagents commonly known in the art.

In all of these assays and methods, a number of distinct peptides or antibodies may be used, either sequentially or simultaneously, and differently labelled, in order to detect a number of different PoERV types in a single assay.

Diagnostic assays based upon the present invention may be used to determine the presence or absence of PoERV infection, and the PoERV type involved.

In an assay for the diagnosis of viral infection, there are basically three distinct approaches that can be adopted, involving the detection of viral nucleic acid, viral antigen or viral antibody respectively. Viral nucleic acid is generally regarded as the best indicator of the presence of the virus itself and would identify materials likely to be infectious. However, the detection of nucleic acid is not usually as straightforward as the detection of antigens or antibodies since the level of target can be very low. Viral antigen is used as a marker for the presence of virus and as an indicator of infectivity. Depending upon the virus, the amount of antigen present in a sample can be very low and difficult to detect. Antibody detection is relatively straightforward because, in effect, the host immune system is amplifying the response to an infection by producing large amounts of circulating antibody. The nature of the antibody response can often be clinically useful, for example IgM rather than IgG class antibodies are indicative of a recent infection, or the response to a particular viral antigen may be associated with clearance of the virus. Thus the exact approach adopted for the diagnosis of a viral infection depends upon the particular circumstances and the information sought. In the case of PoERV, a diagnostic assay may embody any one or a combination of these three approaches.

In an assay for the diagnosis of PoERV involving detection of viral antigen or antibody, the method may comprise contacting a test sample with a peptide of the present invention or a polyclonal or monoclonal antibody
5 against the peptide and determining whether there is any antigen-antibody binding contained within the test sample. For this purpose, a test kit may be provided comprising a peptide, as defined herein, or a polyclonal or monoclonal antibody thereto and means for determining whether there is
10 any binding with antibody or antigen respectively contained in the test sample to produce an immune complex. The test sample may be taken from any appropriate tissue or physiological fluid, such as blood (e.g. serum or plasma), saliva, urine, cerebrospinal fluid, sweat, tears or tissue
15 exudate. If a physiological fluid is obtained, it may optionally be concentrated for any viral antigen or antibody present.

A variety of assay formats may be employed. The peptide can be used to capture selectively antibody against
20 PoERV from solution, to label selectively the antibody already captured, or both to capture and label the antibody. In addition, the peptide may be used in a variety of homogeneous assay formats in which the antibody reactive with the peptide is detected in solution with no
25 separation of phases.

The types of assay in which the peptide is used to capture antibody from solution involve immobilization of the peptide on to a solid surface. This surface should be

capable of being washed in some way. Examples of suitable surfaces include polymers of various types (moulded into microtitre wells; beads; dipsticks of various types; aspiration tips; electrodes; and optical devices),
5 particles (for example latex; stabilized red blood cells; bacterial or fungal cells; spores; gold or other metallic or metal-containing sols; and proteinaceous colloids) with the usual size of the particle being from 0.02 to 5 microns, membranes (for example of nitrocellulose; paper;
10 cellulose acetate; and high porosity/high surface area membranes of an organic or inorganic material).

The attachment of the peptide to the surface can be by passive adsorption from a solution of optimum composition which may include surfactants, solvents, salts and/or chaotropes; or by active chemical bonding. Active bonding
15 may be through a variety of reactive or activatable functional groups which may be exposed on the surface (for example condensing agents; active acid esters, halides and anhydrides; amino, hydroxyl, or carboxyl groups; sulphhydryl
20 groups; carbonyl groups; diazo groups; or unsaturated groups). Optionally, the active bonding may be through a protein (itself attached to the surface passively or through active bonding), such as albumin or casein, to which the viral peptide may be chemically bonded by any of
25 a variety of methods. The use of a protein in this way may confer advantages because of isoelectric point, charge, hydrophilicity or other physico-chemical property. The viral peptide may also be attached to the surface (usually

but not necessarily a membrane) following electrophoretic separation of a reaction mixture, such as immunoprecipitation.

After contacting the surface bearing the peptide with
5 a test sample (in the presence of a blocking mixture if required), allowing time for reaction, and, where necessary, removing the excess of the sample by any of a variety of means, (such as washing, centrifugation, filtration, magnetism or capillary action) the captured
10 antibody is detected by any means which will give a detectable signal. For example, this may be achieved by use of a labelled molecule or particle as described above which will react with the captured antibody or any molecule containing an epitope contained in the peptide. In one
15 embodiment, it is preferred to add an anti-human IgG conjugated to horseradish peroxidase and then to detect the bound enzyme by reaction with a substrate to generate a colour.

The detectable signal may be produced by any means
20 known in the art such as optical or radioactive or physico-chemical and may be provided directly by labelling the molecule or particle with, for example, a dye, radiolabel, fluorescent, luminescent, chemiluminescent, electroactive species, magnetically resonant species or fluorophore, or
25 indirectly by labelling the molecule or particle with an enzyme itself capable of giving rise to a measurable change of any sort. Alternatively the detectable signal may be obtained using, for example, agglutination, or through a

diffraction or birefringent effect if the surface is in the form of particles.

Assays in which a peptide itself is used to label an already captured antibody require some form of labelling of the peptide which will allow it to be detected. The labelling may be direct by chemically or passively attaching for example a radiolabel, magnetic resonant species, particle or enzyme label to the peptide; or indirect by attaching any form of label to a molecule which will itself react with the peptide. The chemistry of bonding a label to the peptide can be directly through a moiety already present in the peptide, such as an amino group, or through an intermediate moiety, such as a maleimide group. Capture of the antibody may be on any of the surfaces already mentioned in any reagent including passive or activated adsorption which will result in specific antibody or immune complexes being bound. In particular, capture of the antibody could be by anti-species or anti-immunoglobulin-sub-type, by rheumatoid factor, proteins A, G and the like, or by any molecule containing an epitope contained in the peptide.

The labelled peptide may be used in a competitive binding fashion in which its binding to any specific molecule on any of the surfaces exemplified above is blocked by antigen in the sample. Alternatively, it may be used in a non-competitive fashion in which antigen in the sample is bound specifically or non-specifically to any of the surfaces above and is also bound to a specific bi- or

poly-valent molecule (e.g. an antibody) with the remaining valencies being used to capture the labelled peptide.

Examples of the invention will now be described by way of illustration only, and with reference to the accompanying Figures, in which:

Figure 1 is the consensus amino acid sequence of the PoERV GAG protein; Figure 2 is a comparison of amino acid sequences of five different PoERV ENV proteins (PERV; PERV A, PERV B, PoEV 1; Galbraith et al., 1997; PERV MSL, Tsukuba; Fishman, 1997) and Figure 3 is a comparison of amino acid sequences of the variable region of five different PoERV ENV proteins, showing the six different ENV peptides (peptides D-H and J) referred to in the following examples.

METHODS

Preparation of PoERV virions.

Human 293 cells (American Type Culture Collection [ATCC] # CRL1573) and Raji cells (ATCC # CCL 86) were infected with PoERV by exposure to polybrene (Sigma-Aldrich Co. Ltd.) and continued incubation with cell-free filtered supernatant from PK-15 (ATCC # CCL 33) cells previously shown to be infected with all three subgroups of PoERV. The 293 cells allow replication of type B PoERV (POEV-1). The 293 cells were shown to be infected after passage by measurement of the reverse transcriptase activity of the cell supernatant and by a PoERV GAG-specific Polymerase Chain Reaction (PCR). The resulting virus particles were isolated from the cell line supernatant as follows.

WO 01/12816

PCT/GB00/03159

15

Supernatant from exponentially growing cells was layered onto a 20/40% (w/v) discontinuous sucrose density gradient and centrifuged at 100,000 g for 150 min. The viral material at the sucrose interface was harvested, and viral particles pelleted by further ultracentrifugation at 100,000g for 60 min, followed by resuspension in DMEM (Life Technologies Ltd., UK).

Control retroviruses

To provide retroviral controls for cross reactivity with PoERV GAG and ENV, Squirrel monkey retrovirus, Murine leukaemia virus, Maedi-Visna virus and Equine infectious anemia virus virions were prepared from the appropriate infected cell line as described by Shepherd and Smith (1999).

Selection and preparation of GAG peptides

Peptides from the GAG protein can provide a capture antigen and a means to generate positive control antisera. The antisera can be directed against conserved polypeptides present in the PoERV virion core likely to induce an immune response in recipients of the virus. These reagents would be useful diagnostic tools for immunosurveillance of recipients of porcine material or tissues for exposure to PoERV. Therefore, peptides encompassing potential antigenic regions of PoERV GAG were selected from the translated amino-acids derived from the sequence of the gag region of PoERV based on three criteria; hydrophilicity,

potential β -turns and K, D, R and E charged residues. The regions were identified using Hopp and Woods hydropathicity (1981) scale and Kyte and Doolittle (1982) hydrophobicity scale.

5 For GAG two peptides were identified as potential antigens. Peptide 1 was from the C-terminus of p30-GAG at residue 437-451 of the polypeptide, nucleotides 1896-1940 of the gag open reading frame (ORF). Peptide 2 was from the start of the p10 segment of the GAG polypeptide at residue
10 502-515 of the polypeptide, nucleotides 2091- 2132 of the gag ORF. The peptides shown below were chemically synthesised by Genosys Biotechnologies Inc.

Peptide 1: (C)REERRDRRQEKNLTK
15 Peptide 2: (W)ARNCPKKGNGPKS

The bracketed amino-acid is not in sequence - 5' position is from next residue (R).

20 These peptides, and the GAG consensus polypeptide sequence, are shown in Figure 1.

A BLAST search (Altschui et al., 1997) of the non-redundant GenBank coding sequences with GAG peptide 1 showed homology with seven sequences all from the gag ORF. Of the seven, three were with PoERV sequences with
25 accessions gi 3116446 (100% match), emb CAA7651 (100% match), gi 3116442 (86% match). The remainder were against the closely related Gibbon ape leukaemia virus (gi 3033415, 92% match) and Simian sarcoma virus (sp PO3330, 86% match).

The remaining two sequences were from murine viruses, including a virus from *Rattus norvegicus* (emb CAA24514; 92% match), and one against *Mus dunni* endogenous retrovirus (gi 3309124, 93% match). A similar BLAST search with GAG peptide 2 showed 100 - 99% homology with only the PoERV sequences listed above.

Selection and Preparation of PoERV ENV peptides

We have previously identified and derived the nucleotide sequence of a unique PoERV type capable of infecting human cells (Galbraith, 1997). Furthermore, it has been shown that the amino acid sequence of the ENV region of various PoERV types contains both conserved and non-conserved regions (Galbraith, 1997; Figure 2). In order to exploit these differences to produce immunological reagents to allow the identification of the type of PoERV giving rise to an immunological reaction in a patient, PoERV-type-specific ENV peptides and antisera were generated.

Six peptides, D-H and J, were identified as potential antigens. The peptides are shown below and their position in the env ORF of the various PoERV types is shown in Figure 3. The peptides were chemically synthesised by the University of Glasgow Veterinary Pathology Department.

Peptide D: TSLRPDITQPPSNSTT
Peptide E: KGKQENIQKWINGMS
Peptide F: RKTGKYSKVDKWEYELGNS
Peptide G: NTVLTGQRPPTQ

WO 01/12816

PCT/GB00/03159

18

Peptide H: GHGRWKDWQQRVQKDVRNKKQIS

Peptide J: IQEQRPSNPNSDYNTT

The amino acid residues of all peptides are identified by
the standard one letter abbreviations.

Preparation of recombinant PoERV p30-GAG and ENV polypeptides

In addition to the peptide reagents more general PoERV p30-GAG and an abbreviated ENV polypeptides were designed and produced for use as capture antigens and to produce anti-polypeptide sera. The required polypeptide portions of the *gag* and *env* genes were produced by PCR amplification, molecularly cloned into a prokaryotic expression vector and expressed as described below using standard techniques (Maniatis et al, 1982).

PoERV p30-GAG

A fragment encompassing the p30 region of the *gag* ORF from nucleotide 1173-1949 of the PoERV genome (Galbraith et al, 1997; Gene Bank Accession # A66553) was amplified by PCR from cDNA generated from PK15 mRNA using ligation independent cloning oligonucleotide primers (pET-32 Ek/LIC cloning and expression vector; Novagen Inc. Catalogue # 69076-3). The oligonucleotides were:

10069082-103002

WO 01/12816

PCT/GB00/03159

19

p30 forward 5' GAC GAC GAC AAG CTG CGC ACC TAT GGC C 3'
 p30 reverse 5' GAG GAG AAG CCC GGG TCT AGG CCA AGA TCT
 TAG TCA AAT TCT TCT C 3'

5 The nucleotides in bold are viral specific.

 The PCR conditions were 30 cycles of 95°C for 1 min,
 58°C for 1 min and 72°C for 1 min. The resulting 776 base
 pair fragment was molecularly cloned into the appropriate
 10 LIC site of the pET-32 LIC vector following the
 manufacturer's instructions (Novagen Inc. 69076-3
 instruction manual), transfected into competent Novoblu[™]
 Escherichia coli cells and plated on solid LB medium
 containing ampicillin. The transformed colonies were
 15 selected by resistance to ampicillin.

PoERV ENV

 A fragment encompassing the region of the env ORF from
 nucleotide 5616- 6304 of the PoERV genome (Galbraith et
 20 al,1997; Gene Bank Accession # A66553) was amplified by PCR
 from cDNA generated from PK15 and PoERV-infected 293 cells
 (PoERV B) mRNA using ligation independent cloning
 oligonucleotide primers (pET-32 Ek/LIC cloning and
 expression vector; Novagen Inc. Catalogue # 69076-3). The
 25 oligonucleotides were:

env forward 5' GAC GAC GAC AAG ATC CAT GCA TCC CAC GTT
 3'

WO 01/12816

PCT/GB00/03159

20

env reverse 5' GAG GAG AAG CCC GGT CTC TAT CCT AAG GCG
 3'

The nucleotides in bold are viral specific.

5

The PCR conditions were 30 cycles of 95°C for 1 min, 60°C for 1 min and 72°C for 1 min. The resulting 688 base pair fragment was molecularly cloned into the appropriate LIC site of the pET-32 LIC vector following the manufacturer's instructions as described above.

For expression from the T7 promoter the recombinants are required to be transferred to a host with T7 polymerase activity. To this end plasmid DNA was isolated from the ampicillin resistant Novoblu[™] clones carrying the gag or env fragment in the correct orientation for expression as determined by restriction endonuclease mapping. The plasmid DNAs were each transfected into competent *E. coli* AD494 (DE3) *trx B*-.

For screening for the production of recombinant protein, two ml cultures of *E. coli* transformed with either of the two expression constructs were grown with shaking at 37°C to late log phase (O.D._{600nm} of approximately 0.6) and induced by the addition of Isopropylthio-beta-galactoside (IPTG) to 0.1 mM. Induced cultures were then incubated for a further 2 h after which the bacteria were collected by centrifugation. The bacterial pellet was lysed by boiling in SDS-PAGE sample buffer and the protein profile of the induced bacteria was analysed on a 12% acrylamide gel

WO 01/12816

PCT/GB00/03159

21

followed by staining with coomasie brilliant blue dye.
(Laemmli, 1970; Gallagher, 1997).

Large scale preparations of purified GAG and ENV
polypeptides were made according to the manufacturer's
5 instructions (Novagen Inc. Catalogue # 69076-3).

Preparation of antisera to whole virions, peptides and
recombinant P30-GAG and ENV antigens.

For GAG peptide 1 and peptide 2, the peptides were
10 conjugated with keyhole limpet hemacyanin carrier protein
and each of two rabbits was inoculated six times at
fourteen day intervals. The animals were bled out at day
seventy seven after the first inoculation. The p30-GAG
polypeptide was inoculated three times at fourteen day
15 intervals into a rabbit. The animal was bled out at day
seventy seven.

For ENV peptides D-H and J, the peptides were
conjugated with keyhole limpet hemacyanin carrier protein
and one sheep was inoculated three times at twenty eight
20 day intervals. The ENV polypeptide was inoculated three
times at fourteen day intervals into a rabbit.

Virions purified from PK15 cells were inoculated three
times at fourteen day intervals into each of two guinea
pigs.

25

followed by staining with coomassie brilliant blue dye.
(Laemmli, 1970; Gallagher, 1997).

Large scale preparations of purified GAG and ENV
polypeptides were made according to the manufacturer's
5 instructions (Novagen Inc. Catalogue # 69076-3).

Preparation of antisera to whole virions, peptides and
recombinant P30-GAG and ENV antigens.

For GAG peptide 1 and peptide 2, the peptides were
10 conjugated with keyhole limpet hemacyanin carrier protein
and each of two rabbits was inoculated six times at
fourteen day intervals. The animals were bled out at day
seventy seven after the first inoculation. The p30-GAG
polypeptide was inoculated three times at fourteen day
15 intervals into a rabbit. The animal was bled out at day
seventy seven.

For ENV peptides D-H and J, the peptides were
conjugated with keyhole limpet hemacyanin carrier protein
and one sheep was inoculated three times at twenty eight
20 day intervals. The ENV polypeptide was inoculated three
times at fourteen day intervals into a rabbit.

Virions purified from PK15 cells were inoculated three
times at fourteen day intervals into each of two guinea
pigs.

Indirect immunofluorescence staining

To test the specificity of the p30-GAG antisera the method outlined by Riggs (1989) was used. PoERV-infected Raji cells and uninfected control Raji cells were fixed and tested for indirect immunofluorescence with anti p30-GAG using a fluorescein isothiocyanate (FITC) labelled anti-rabbit detector antibody. The cells were examined by fluorescence microscopy.

Preparation of Western blot membranes

Recombinant p30-GAG polypeptide and ENV polypeptide were prepared, harvested and purified from an *E.coli* vector. The recombinant proteins were tested to determine an appropriate dilution of protein which yielded a positive result in the immunoassay. In addition, extracts from PoERV-infected 293 cells, PoERV-infected Raji cells or purified PoERV virions were used as antigens. To obtain specific and reproducible Western blot assays, a number of parameters were required to be optimised for each assay, such as: Primary antibody dilution, incubation time, incubation temperature, secondary antibody dilution, incubation time, incubation temperature, washing buffers, blocking/dilution buffers, developing reagents. Recombinant polypeptides were added to nine wells of a ten lane 12% Tris/glycine acrylamide gel. Molecular weight markers were added to the first lane. The samples were electrophoresed and the gel electroblotted to a polyvinylidene fluoride (PVDF) membrane. (Gallagher et al,

WO 01/12816

PCT/GB00/03159

23

1997). The membrane was cut into strips each strip containing one lane of recombinant protein. These strips were used as the basis of the assay.

5 **Preparation dilutions of antisera**

Samples were prepared in a Class 2 safety cabinet or other clean environments.

10 A typical negative control was prepared by making up to a 1:200 dilution of normal sera in blocking reagent.

15 A typical positive control was prepared by making a 1: 500, 1: 1000 or greater dilution of anti-PoERV p30-GAG polypeptide, peptide serum or anti - recombinant ENV serum.

20 A typical test serum was prepared by making up to a 1:200 dilution of sera.

25 **Preparation of Western blotting membranes and PoERV antibody detection.**

To block non-specific binding sites membrane strips each were placed in a 15 ml centrifuge tube and 2 ml blocking reagent (2.5 g skimmed dried milk in 50 ml PBS/ 0.5% v/v Tween-20TM) added. The strips were placed on a rotary shaker such that the strip moved slightly on each revolution and were incubated for 30 min at ambient

temperature. The blocking reagent was removed and replaced with 5-10 μ l of the diluted serum. The membrane was incubated with shaking for 1 h at ambient temperature. To stop incubation the strip was removed from diluted serum and placed into PBS/ Tween-20™ and washed with three changes of PBS/ Tween-20™ at ambient temperature with shaking.

The appropriate species specific secondary antiserum conjugated to alkaline phosphatase was used as detector e.g. if human serum was being tested, an anti-human IgG alkaline phosphatase (AP) conjugate was used. The p30-GAG positive control required anti-rabbit IgG AP conjugate for detection and the anti ENV required anti-sheep IgG AP conjugate. The detection was done as follows; each strip was placed in an unused 15 ml centrifuge tube, 2 ml of 1:1000 dilution of secondary sera in blocking reagent was added and incubated with shaking at ambient temperature for 1 h. The strip was removed from the centrifuge tube, placed in PBS/Tween-20™ and washed with 3 changes of PBS/Tween-20™, at ambient temperature with shaking. The strips were then put into a 15 ml centrifuge tube and 2 ml of bromochlorindoyl phosphate/ nitroblue tetrazolium (BCIP/NBT, Sigma-Aldrich Co. Ltd.) solution was added to each tube. The strips were shaken gently and allowed to develop for 5 min. The reaction was stopped by rinsing the membrane strip in purified water and the strips were removed from the water and allowed to air dry.

ENZYME-LINKED IMMUNOSORBENT ASSAY (ELISA)**Antigen coating of microtiter plates**

50 μ l of antigen (either recombinant p30-GAG, extracts from PoERV-infected 293 cells or purified PoERV B virions) were diluted to the required concentration in carbonate-bicarbonate coating buffer (Sigma-Aldrich Co. Ltd.) was added to wells of a 96 well, flat bottomed microtiter ELISA plate (Dynex Immulon 2). Some wells contained carbonate-bicarbonate coating buffer only and some were left blank to act as controls for non-specific binding. The plate was covered with a plate seal and incubated at 4°C for approximately 16 h. The unbound antigen and coating solution were then removed from the wells with a pipette and washed three times with PBS / 0.05% Tween-20™. Any remaining PBS / 0.05% Tween-20™ was removed by blotting on a tissue.

Blocking of microtiter plates

50 μ l of fresh blocking buffer (5% (w/v) skimmed milk / PBS / 0.05% Tween-20™) was added to each antigen coated well and control well. The plate was covered with a plate seal and incubated at ambient temperature in an humidified chamber for 1 h. The blocking buffer was removed and the plate wells washed three times with PBS / 0.05% Tween-20™ and any remaining PBS / 0.05% Tween-20™ was removed by blotting on a tissue.

Incubation with primary antibody

50 μ l of negative control sera, test sera, and positive control sera at the experimental dilution were added to the antigen coated wells and the plates covered with a plate seal and incubated at ambient temperature for approximately 1 h in an humidified chamber. Following incubation all sera were removed using a pipette and the plate wells washed three times with PBS / 0.05% Tween-20™, any remaining PBS / 0.05% Tween-20™ was removed by blotting on a tissue.

Incubation and development with peroxidase-conjugated secondary antibody

50 μ l of a 1:500 dilution of species specific peroxidase conjugated secondary antibody in blocking buffer was added to each antigen coated well. For human serum an anti-human IgG peroxidase conjugate was used. The p30-GAG positive control required anti-rabbit IgG peroxidase conjugate for detection and the anti ENV required anti-sheep IgG peroxidase conjugate. The plates were covered with a plate seal and incubated at ambient temperature for approximately 1 h in an humidified chamber. Following incubation the conjugate was removed using a pipette and the plate wells washed three times with PBS / 0.05% Tween-20™. Any remaining PBS / 0.05% Tween-20™ was removed by blotting on a tissue.

The substrate was prepared as follows: one O-phenylenediamine tablet (Sigma-Aldrich Co. Ltd.) and one urea/H₂O₂ tablet were dissolved in 20 ml of purified water.

An aliquot of 50 μ l of substrate was added to each well and the plate incubated at ambient temperature in the dark for 30 min. The reaction was then stopped by adding 50 μ l of 3N HCl or 3 M H₂SO₄ to each well. The colour development in the wells was measured at 490 nm using a Dynex MRX microplate reader.

For alkaline phosphatase conjugated secondary antibodies the substrate used was p-Nitrophenyl phosphate (pNPP; Sigma-Aldrich Co. Ltd.) and the plates were read at 405 nm.

Electron Microscopy

Negative stain electron microscopy (NSEM) (Doane, 1980) was used to identify the presence of PoERV virions. Supernatant from PoERV-infected pK15 cells was layered onto a 20/40% (w/v) discontinuous sucrose density gradient and centrifuged at 100,000 g for 150 min. The viral material at the sucrose interface was harvested, and viral particles pelleted by further ultracentrifugation at 100,000g for 60 min, followed by resuspension in DMEM (Life Technologies Ltd.). The sample was then applied to pioloform-coated copper 300 mesh EM grids and allowed to air dry. Grids were fixed with 2.5% glutaraldehyde (Agar Scientific), stained with 5% uranyl acetate (Agar Scientific) and allowed to air dry. Grids were examined on a Philips EM-400 transmission electron microscope.

WO 01/12816

PCT/GB00/03159

28

Immuno- Electron Microscopy

The immunostaining was done following the method of Marshall et al (1992). Briefly, NSEM samples were applied to pioloform-coated nickel 300 mesh EM grids and allowed to
5 air dry. Grids were fixed with modified immunofix, post fixed with 0.5 M NH_4Cl , then incubated with 2% bovine serum albumin (Sigma-Aldrich Co. Ltd.). Samples were then incubated with rabbit anti-PoERV (rabbits immunised with whole PoERV) or rabbit anti-PoERV p30-GAG antibody, washed
10 in modified immunobuffer followed by incubation with anti-rabbit IgG gold conjugate (Sigma-Aldrich Co. Ltd.). Grids were stained with 5% uranyl acetate, and allowed to air dry. Samples were visualised on a Philips EM-400 transmission electron microscope.

15

EXAMPLE ONE

Indirect immunofluorescence staining of PoERV-infected cells.
20 Viral specific fluorescence was observed in PoERV-infected Raji cells using the anti p30-GAG antiserum. No immunofluorescence was seen with the negative control Raji cells.

25

WO 01/12816

PCT/GB00/03159

29

EXAMPLE TWO**Western Blotting for Antibodies to PoERV GAG**

Anti-GAG peptide 1 antisera and sera from rabbits
5 inoculated with the recombinant p30-GAG polypeptide
detected the expected protein of approximately 30kd in
extracts of PoERV-infected 293 or PoERV-infected Raji
cells, purified PoERV virions and recombinant p30-GAG. The
PoERV antibody could be detected at a dilution of 1:1000.

10

No band of equivalent size to the GAG 30 kd polypeptide was
detected in uninfected control cells.

No band of equivalent size to the GAG 30 kd polypeptide was
15 detected against the following purified retroviruses:

Squirrel monkey retrovirus
Murine leukaemia virus
Maedi-Visna virus
20 Equine infectious anemia virus

Therefore, the positive control antisera were specific for
PoERV.

25 Determination of assay specificity using Serum Panels**Normal human sera**

On testing of 90 normal serum samples from healthy
individuals, whose blood was taken for occupational health

WO 01/12816

PCT/GB00/03159

30

reasons, no PoERV reactive antibody was detected in any of the sera.

Normal primate sera

On testing 42 normal serum samples from healthy primates no
5 PoERV reactive antibody was detected. There was no cross-reactivity with sera from normal primates.

Cardiac transplant patient sera

On testing 20 serum samples from individuals who had
10 received a cardiac transplant in the preceding 36 months no PoERV reactive antibody was detected. There was no cross-reactivity with sera from patients who had been immunosuppressed.

HIV positive sera

15 On testing 13 serum samples from individuals positive for the presence of antibody to HIV no PoERV reactive antibody was detected. There was no cross-reactivity with sera from patients infected with a human Lentivirus.

20 HTLV positive sera

On testing 10 serum samples from individuals positive for the presence of antibody to HTLV-1 virus no PoERV reactive antibody was detected. There was no cross-reactivity with sera from patients infected with a human Gammaretrovirus.

25

WO 01/12816

PCT/GB00/03159

31

Butchers with acute lymphoblastic leukaemia (ALL)

On testing 3 serum samples from butchers with ALL no PoERV reactive antibody was detected. There was no cross-reactivity with sera from patients with assumed prolonged exposure to PoERV and PoERV antigens with a leukaemic disorder.

EXAMPLE THREE**ELISA p30-GAG**

A titration of p30-GAG antigen to anti-p30-GAG antisera gave a significant signal at 1:250600 dilution antigen to 1:32000 dilution of antisera. A similar titration of antisera against PoERV virions gave a significant signal at a 1:3200 dilution of both antigen and antisera.

Normal human sera

On testing of five normal serum samples from healthy individuals whose blood was taken for occupational health reasons, no significant signal was detected in any of the sera against recombinant p30-GAG.

EXAMPLE FOUR**Detection and Visualisation of PoERV Virions by Immunoelectron Microscopy**

Examination of PoERV virion preparations by negative stain revealed particles showing the characteristic size and structure of a Gammaretrovirus of approximately 90-120 nm with a dark inner core and double membranous outer region.

The particles bound immuno-gold labeled anti p30-GAG antiserum indicating that the antiserum could be used to visualise PoERV virions by immuno-electron microscopy.

5 **EXAMPLE FIVE**

ELISA ENV

A titration of antisera raised against ENV peptides D, E, F, G, H, J. to purified PoERV B virions gave a significant signal indicating that the ENV peptides produced a virus-specific reaction in the animals. Peptides D and F, both
10 from PoERV B (POEV1; Figure 2; Galbraith et al, 1997) gave the highest signal.

EXAMPLE SIX

15 **Western Blotting for Antibodies to PoERV ENV**

Antisera from guinea pigs inoculated with whole purified PoERV virions from PK15 cells detected the expected recombinant ENV protein of approximately 24 kD in extracts of *E.coli* expressing the *env* construct.

20 No band of equivalent size to the ENV 24 kD polypeptide was detected on *E.coli* control cells without the expression construct.

REFERENCES

- Altschui SF, Madden TL, Schaffer AA, Zhang J, Zhang Z,
Miller W and Lipman DJ (1997). Gapped BLAST and PSI-BLAST:
5 a new generation of protein database search programs.
Nucleic Acids Research, 25: 3389-3402.
- Doane FW . (1980). Virus morphology as an aid for rapid
diagnosis. Yale Journal of Biological Medicine, 53, 19-25.
- 10 Ekins RP. (1997). Immunoassay design and optimisation. In
Principles and Practice of Immunoassay Second Edition. Ed.
CP Price and DJ Newman. Macmillan Reference, Ltd. London.
- 15 Fishman JA. (1997). Patent Application WO97/21836.
- Galbraith DN. Haworth C, Lees GM, and Smith KT. (1997).
Patent Application WO97/40167.
- 20 Gallagher S. (1997). One-dimensional SDS gel
electrophoresis of proteins. In Current Protocols in
Molecular Biology. 10.2.2-10.2.35. John Wiley & Sons, Inc.
USA.
- 25 Gallagher S, Winston SE, Fuller SA and Hurrell JGR. (1997).
Immunoblotting and immunodetection. In Current Protocols in
Molecular Biology. 10.8.1-10.8.21. John Wiley & Sons, Inc.
USA.

- Hopp TP and Woods KR. (1981). Prediction of protein antigenic determinants from amino acid sequences. Proceeding of the National Academy of Sciences USA, 78, 3824-3828.
- 5 Kyte J and Doolittle RF. (1982). A simple method for displaying the hydropathic character of a protein. Journal of Molecular Biology, 157, 105-132.
- 10 Laemmli UK. 1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. Nature, 227, 680-685.
- 15 Maniatis T, Fritsch EF and Sambrook J. (1982). Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- Marshall GE, Kontas AG, Reid GG, Edwards JG and Lee WR. (1992). Type IV collagen and laminin in Bruch's membrane and basal linear deposit in the human macula. British
- 20 Journal of Ophthalmology, 76, 607-614.
- Onions D, Hart D, Mahoney C, Galbraith D and Smith K. (1998). Endogenous retroviruses and the safety of porcine xenotransplantation. Trends in Microbiology, 6, 430-431.
- 25 Patience C, Takeuchi Y and Weiss R. (1997). Infection of human cells by an endogenous virus of pigs. Nature

Medicine, 3, 282-286.

5 Riggs JL. (1989). Immunofluorescence staining. In
Diagnostic Procedures for Viral, Rickettsial and Chlamydial
Infections Sixth Edition. Ed. NJ Schmidt and RW Emmons.
123-133. American Public Health Association Inc. Washington
DC.

10 Strandstrom H, Veijalainen P, Moennig V, Hunsmann G,
Schwarz H and Schafer W. (1974). C-type particles produced
by a permanent cell line from a leukemic pig. 1. Origin and
properties of the host cells and some evidence for the
occurrence of C-type particles. Virology, 57, 175-178.

15 Shepherd AJ and Smith KT. (1999). Viral evaluation of
animal cell lines used in biotechnology. In Methods in
Biotechnology, Vol. 8: Animal Cell Biotechnology. Ed. N
Jenkins. Humana Press, Inc. Totwa New Jersey.

20 Todaro GJ, Benveniste RE, Lieber MM, and Sher CJ. (1974).
Characterisation of a type C virus released from the
porcine cell line PK (15). Virology, 58, 65-74.

CLAIMS

1. A porcine endogenous retrovirus (PoERV) polypeptide fragment, wherein said polypeptide fragment has PoERV specific antigenic or immunogenic activity.
- 5 2. A polypeptide fragment according to claim 1, wherein said polypeptide fragment comprises a fragment of a PoERV GAG polypeptide.
- 10 3. A polypeptide fragment according to claim 2, wherein said polypeptide fragment comprises a fragment of a polypeptide sequence as shown in Figure 1.
- 15 4. A polypeptide fragment according to claim 3, wherein said polypeptide fragment comprises the amino acid sequence REERRDRRQEKNLTK.
- 20 5. A polypeptide fragment according to claim 3, wherein said polypeptide fragment comprises the amino acid sequence ARNCPKKGNKGPKV.
- 25 6. A polypeptide fragment according to claim 1, wherein said polypeptide fragment comprises a fragment of a PoERV ENV polypeptide.
7. A polypeptide fragment according to claim 6 wherein said polypeptide fragment comprises a fragment of a polypeptide sequence as shown in Figures 2 or 3.

8. A polypeptide fragment according to claim 7 wherein said polypeptide fragment comprises a fragment from within a conserved region of the sequences shown in Figures 2 and 3.

5

9. A polypeptide fragment according to claim 7 wherein said polypeptide fragment comprises a fragment from within a non-conserved region of the sequences shown in Figures 2 and 3.

10

10. A polypeptide fragment according to claim 9 wherein said polypeptide fragment comprises an amino acid sequence selected from the sequences of peptides D, E, F, G, H and J as shown in Figure 3.

15

11. A polypeptide fragment according to claim 7 wherein said polypeptide fragment comprises a fragment from within a conserved region of the sequences shown in Figures 2 and 3, and a fragment from within a non-conserved region of the sequences shown in Figures 2 and 3.

20

12. A polypeptide fragment according to claim 1, wherein said polypeptide fragment comprises a fragment of a PoERV GAG polypeptide and a fragment of a PoERV ENV polypeptide.

25

13. An antiserum specific to a polypeptide fragment in accordance with any preceding claim.

WO 01/12816

PCT/GB00/03159

38

14. A PoERV specific antibody or fragment thereof raised against a polypeptide fragment in accordance with any of claims 1 to 12.

5 15. Use of an antibody according to claim 14 in the detection of PoERV in a sample.

10 16. Use of a polypeptide fragment according to any one of claims 1 to 12 in the detection of PoERV antibodies in a sample.

15 17. An assay kit for use in detection of PoERV in a sample, the kit comprising an antibody in accordance with claim 14.

18. An assay kit for use in detection of PoERV antibodies in a sample, the kit comprising a polypeptide fragment in accordance with any of claims 1 to 12.

20 19. Use of an antibody according to claim 14, or a polypeptide fragment according to any one of claims 1 to 12 in therapy or diagnosis.

25

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
22 February 2001 (22.02.2001)

PCT

(10) International Publication Number
WO 01/12816 A2(51) International Patent Classification: C12N 15/49,
C07K 14/15, G01N 33/50, C07K 16/10, A61K 39/21(74) Agents: MCCALLUM, William, Potter et al.: Cruik-
shank & Fairweather, 19 Royal Exchange Square, Glasgow
G1 3AE (GB).

(21) International Application Number: PCT/GB00/03159

(22) International Filing Date: 16 August 2000 (16.08.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9919604.0 18 August 1999 (18.08.1999) GB(71) Applicant (for all designated States except US): QUIP
TECHNOLOGY LIMITED [GB/GB]; Todd Campus,
West of Scotland Science Park, Glasgow G20 0XA (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): GALBRAITH,
Daniel [GB/GB]; 513 Alexandra Parade, Glasgow G31
3EW (GB). KELLY, Helena [GB/GB]; Flat 1/2, 19
Kelvindale Gardens, Glasgow G20 8DW (GB). SMITH,
Kenneth [GB/GB]; 5 Capelrig Lane, Newton Mearns,
Glasgow G77 6XZ (GB).

Published:

— Without international search report and to be republished
upon receipt of that report.For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: RETROVIRUS ASSAY

PoERV GAG POLYPEPTIDE SHOWING POSITIONS OF PEPTIDES

MGQIVTITPLSLTDHWTEVRSRAHNLVQVQKGPWQTFCAENWPTFDVWGPSEGTFNSEIILAVKAIIPQ
 TGPQSHPDQEPYIITWQDLAEDPPVWPKPWLKPRKPGPRILALGEKNKSAEKVEPSSSYLPRDRGAAD
 LAGTPTCSPTPLSSTGCCGEGTSAPFGAPVVEGPAAGTSSRRGATPERTDEIAILFLRTYGVPMFGQLOQP
 LQYWPSSADLYNMKTNHPPFSEDPQRLTGLVESLMFSHQPTWDDCQQLQTLFTTEERERILLEARKNV
 PGADGPTQLQNEIDMGFFLTRPGWDYNTAEGRESLKITYRQALVAGLRGASRRPTNLAKVREVMQGNPEP
 PSVFLERLMEAFRRFTFPDPTSEAQKASVALAFIQSALDIRKKQLRLEGQAEALDLVREAKVYVYRR
 ETEEEKRQKEKEREERERRRRQKLNLTkILAAVVEGKSSRRERDFRKIRSGPRQSGNLGNRTPLDK
 DQCAVCKEKGHWARNCPKKGKGNKPKVLALEEDKD

The gag peptides 1 and 2 are shown in lower case bold. Peptide 1:
 REERRRRQKLNLTk; Peptide 2: ARNCPKKGKGNKPKV.

(57) Abstract: The present invention relates to polypeptide fragments derived from porcine endogenous retrovirus (PoERV) GAG and ENV polypeptides, and to their use in detection of PoERV antibodies in a test sample. Also provided are antibodies to GAG and ENV polypeptides, which may be used to detect PoERV in a sample. Polypeptide sequences are provided which are common to several strains of PoERV, as are sequences specific to a single PoERV strain.

1/4

MGQTVTTPLSLTLDHWTEVRSRAHNLSVQVKKGPMQTFCASEWTFDVGWPSEGTFNSEIILAVKAIIFQ
TGPGSHPDQEPYILTWQDLAEDPPPWWKPNLKPGRKPGPRILALGEKNKHSAEKVEPSSSYLPRDRGAAD
LAGTPTCSPTPLSSTGCCCGTSAPPGAPVVEGPAAGTRSRGATPERTDEIAILPLRTYGPPMPGGQLQP
LQYWPFSADLYNWKTNHPPFSDEDPQRLTGLVESLMFESHQPTWDDCQQLQTLTFTTEERERILLEARKNV
PGADGRPTQLQNEIDMGFPPLTRPGWDYNTAEGRESLKITYRQALVAGLRGASRRPTNLAKVREVMQGPNEP
PSVFLERLMEAFRRFTPDPPTSEAKASVALAFIGQSALDIRKKLQRLGLQEAELRDLVREAEKVYYRR
ETEEEEKRKEKEREEREerrdrreknltkILAAVVEGKSSRERERDFRKIRSGFRQSGNLGNRTPLDK
DQCAYCKEKGHWarncpkgnkgpkvLAEEDKD

FIGURE 1

10/069082

2/4

PERVA MHP T L S R R H L P I R G G K P K R L K I P L S F A S I A W F L T L S I T P Q V N G K R L V D S P N S H K P L S L T W
POEVMSL M H P T L N R R H L P I R G G K P K R L K I P L S F A S I A W F L T L S I T S Q T N G M R I G D S L N S H K P L S L T W
TSUKUBA M H P T L S R R H L P I R G G K P K R L K I P L S F A S I A W F L T L S I T S Q T N G M R I G D S L N S H K P L S L T W
PERVB M H P T L S W R H L P T R G G E P K R L R I P L S F A S I A W F L T L T I T P Q A S S K R L I D S S N P H R P L S L T W
POEV1 M H P T L S R R H L P T R G G E P K R L R I P L S F A S I A W F L T L T I T P Q A S S K R L I D S S N P H R P L S L T W
***** * * * * : * : * * * : * * * * * : * * * * * : * * * * * : * * * * *

PERVA L L T D S G T G I N I N N T Q G E A P L G T W W P E L Y V C L R S V I P S L --- T S P P D I L H A H G F Y V C P G P P
POEVMSL L I T D S G T G I N I N N T Q G E A P L G T W W P D L Y V C L R S V I P S L --- T S P P D I L H A H G F Y V C P G P P
TSUKUBA L I T D S G T G I N I N N T Q G E A P L G T W W P D L Y V C L R S V I P S L --- T S P P D I L H A H G F Y V C P G P P
PERVB L I I D P D T G V T V N S T R G V A P R G T W W P E L H F C L R L I N P A V K --- S T P P N L V R S Y G F Y C C P G - T
POEV1 L I I D P D T G V T V N S T R G V A P R G T W W P E L H F C L R L I N P A V K --- S T P P N L V R S Y G F Y C C P G - T
* : *

PERVA N N E E Y C G N P O D F F C K Q W S C I T S N D G N K W P V S Q D R V S Y S F V N N P T S Y N Q E N Y H G H R K W D
POEVMSL N N G K H C G N P R D F F C K Q W N C V T S N D G Y W K W P T S Q D R V S F S Y V N T Y T S S G Q F N Y --- L T
TSUKUBA N N G K H C G N P R D F F C K Q W N C V T S N D G Y W K W P T S Q D R V S F S Y V N T Y T S S G Q F N Y --- L T
PERVB E K E Y C G G S G E S F C R R W S C V T S N D G D K W P I S Q D R V K F S F V N S G --- P G K Y K V M K ---
POEV1 E K E Y C G G S G E S F C R R W S C V T S N D G D K W P I S Q D R V K F S F V N S G --- P G K Y K M K M ---
: : *

PERVA W Q Q R V Q K D V R N K Q I S C H S L D L D Y L K I S F T E --- K G Q E N I Q K W N G I S W G I V Y Y G G S G R K K
POEVMSL W --- L T G S P K C S P S D L D Y L K I S F T E --- K G Q E N I L K W N G M S W G M V Y Y G G S G K O P
TSUKUBA W --- L T G S P K C S P S D L D Y L K I S F T E --- K G Q E N I L K W N G M S W G M V Y Y G G S G K O P
PERVB W --- L Y K D K S C S P S D L D Y L K I S F T E --- K G Q E N I Q K W N G I S W G I V Y Y K Y G G G - A
POEV1 W --- L Y K D K S C S P S D L D Y L K I S F T E R K T G K Y S K V D K W Y --- E L G N S F L Y G G G - A
* . * * * * * * * . * : *

PERVA G S V L T I R L R I E T Q M E P P V A I G P N K G L A E Q G P P I Q E Q R P - S P N P S D --- Y N T T
POEVMSL G S I L T I R L K I N - Q L E P P M A I G P N T V L T G Q R P P T Q --- G P G P S --- S N I T
TSUKUBA G S I L T I R L K I N - Q L E P P M A I G P N T V L T G Q R P P T Q --- G P G P S --- S N I T
PERVB G S T L T I R L R I E T G E P P V A V G P D K V L A E Q G P P A L E P P H N L P V P Q L T S L R P D I T Q P P S N G T
POEV1 G S T L T I R L R I E T G E P P V A M G P D K V L A E Q G P P A L E P P H N L P V P Q L T S L R P D I T Q P P S N S T
* * * * * : * : * * * : * : * : * : * : * : * : * : * : * : * : * : * : * : * : * : * : * : *

PERVA G S G V P T --- E P N I T I K T G A K L F S L I Q A F O A L N S T T P E A T S S C W L C L A S G P P Y Y E G M A
POEVMSL G S G S D P T --- E S N S T T K M G A K L F S L I Q A F O A L N S T T P E A T S S C W L C L A S G P P Y Y E G M A
TSUKUBA G S G S D P T --- E S S T T K M G A K L F S L I Q A F O A L N S T T P E A T S S C W L C L A S G P P Y Y E G M A
PERVB T G L I P T N T P R N S P G V P V K T G O R L F S L I Q A F O A I N S T D P D A T S S C W L C L S S G P P Y Y E G M A
POEV1 T G L I P T N T P R N S P G V P V K T G O R L F S L I Q A F O A I N S T D P D A T S S C W L C L S S G P P Y Y E G M A
: : *

PERVA R G G K F N V T K E H R D Q C T W G S Q N K L T L T E V S G K G T C I G K V P P S H Q H L C N H T E A F N R T S E S Q Y
POEVMSL R R G K F N V T K E H R D Q C T W G S Q N K L T L T E V S G K G T C I G K V P P S H Q H L C N H T E A F N T S E S Q Y
TSUKUBA R R G K F N V T K E H R D Q C T W G S Q N K L T L T E V S G K G T C I G K V P P S H Q H L C N H T E A F N T S E S Q Y
PERVB K E G K F N V T K E H R N Q C T W G S R N K L T L T E V S G K G T C I G K A P P S H Q H L C Y S T V V Y E Q A S E N Q Y
POEV1 K E R K F N V T K E H R N Q C T W G S R N K L T L T E V S G K G T C I G K A P P S H Q H L C Y S T V V Y E Q A S E N Q Y
: * * * * * : * : * * * : * : * * * : * : * * * : * : * * * : * : * * * : * : * * * : * : * * * : * : * * *

PERVA L V P G Y D R W W A C N T G L T P C V S T L V F N Q T K D F C V M Q I V P R V Y Y P E K A V L D E Y D Y R N R P K
POEVMSL L V P G Y D R W W A C N T G L T P C V S T L V F N Q T K D F C I M Q I V P R V Y Y P E K A I L D E Y D Y R N R H R Q K
TSUKUBA L V P G Y D R W W A C N T G L T P C V S T L V F N Q T K D F C I M Q I V P R V Y Y P E K A I L D E Y D Y R N R H R Q K
PERVB L V P G Y N R W W A C N T G L T P C V S T S V F N Q S K D F C V M Q I V P R V Y Y H P E E V L D E Y D Y R N R P K
POEV1 L V P G Y N R W W A C N T G L T P C V S T S V F N Q S K D F C V M Q I V P R V Y Y H P E E V L D E Y D Y R N R P K
* * * * * : * : * * * : * : * * * : * : * * * : * : * * * : * : * * * : * : * * * : * : * * *

25.10.2000

10/069082

3/4

```
PERVA      REPISLTAVMLGLGVAAGVGTGAALITGPQPLEKGLSNLHRIVTEDLQALESVSNLE
POEVMSL    REPISLTAVMLGLGVAAGVGTGAALTVPGPQLETGLSNLHRIVTEDLQALESVSNLE
TSUKUBA    REPISLTAVMLGLGVAAGVGTGAALTVPGLTGLSNLHRIVTEDLQALESVSNLE
PERVB      REPVSLTLAVMLGLGTAVGVGTGAALITGPQPLEKGELGHAAMTEDLRALAESVSNLE
POEV1      REPVSLTLAVMLGLGTAVGVGTGAALTITGPQPLEKGLGELGHAAMTEDLRALKESVSNLE
*:*****:          *:*****:*****:*****:*****:*****:*****:
```

[illegible]

PERVA REREADQGWFEQWNRSPMWTLLSALTGPLVLVLLLLTVGPCLINRFVAFVRERSAVQ
POEVMSL REKETTTQGWFEGWNRSLWLATLLSALTGPLIVLLLLTVGPCIINKLIAIFIRERISAVQ
TSUKUBA REKETTTQGWFEQWNRSPWLATLLSALTGPLVLVLLLLTVGPCIINKLIAIFIRERISAVQ
PERVB REREADQGWFEQWNRSPMWTLLSALTGPLVLVLLLLTVGPCLINRFVAFVRERSAVQ
POEV1 REREADQGWFEQWNRSPMWTLLSALTGPLVLVLLLLTVGPCLINRFVAFVRERSAVQ

: ***: ***: *: *****:*****:****:***:***:***:***:

```
PERVA      IMVLRQQYQGLLSQGETDL
POEVMSL    IMVLRQQYQSPSSR-EAGR
TSUKUBA    IMVLRQQYQSPSSR-EAGR
PERVB      IMVLRQQYQGLLSQGETDL
POEV1      IMVLRQQYQGLLSQGETDL
*****
*****
```

FIGURE 2

10/069082

4/4

PERVA NNEEYCGNPQDFCFCKQWSCITSNDGWNKWPVSQQDRVSYSFVNNPNTSYNQFNYghgrwkd
 POEVMSL NNGKHCGNPRDFCFCKQWNCVTSNDGYWKWPTSQQDRVSFSYVNTYTSSGQFNY-----LT
 TSUKUBA NNGKHCGNPRDFCFCKQWNCVTSNDGYWKWPTSQQDRVSFSYVNTYTSSGQFNY-----LT
 PERVB EKEKYCGGSGESFCRRWSCVTSNDGDWKWPI SLQDRVKFSFVNSG--PGKYKVMK-----
 POEV1 EKEKYCGGSGESFCRRWSCVTSNDGDWKWPI SLQDRVKFSFVNSG--PGKYKVMK-----

PERVA wqqrqvkdvrnkqisCHSLDLDLKISFTE--KGKQENIQKWNVNGISWGI VYVYGGSGRKK
 POEVMSL W-----IRTGSPKCSPSDDL DYLKISFTE--KGKQENILKWNVNGMSWGMVYVYGGSGKQP
 TSUKUBA W-----IRTGSPKCSPSDDL DYLKISFTE--KGKQENILKWNVNGMSWGMVYVYGGSGKQP
 PERVB -----LYKDKSCSPSDDL DYLKISFTE--kgkqeniqtwingmsWGI VFYKYGGG-A
 POEV1 -----LYKDKSCSPSDDL DYLKISFTErktgkyskvdwy---elgnsFLLYGGG-A

PERVA GSVLTIRLR IETQMEPPVAIGPNKGLAEQGPPiqeqrp-spnpsd-----yntt
 POEVMSL GSILTI RLKIN-QLEPPMAIGPNTVLTGQRPTQ-----GPGPS-----SNIT
 TSUKUBA GSILTI RLKIN-QLEPPMAIGPntvltgqrptq-----GPGPS-----SNIT
 PERVB GSTLTIRLR IETGTEPPVAVGPKVLAEQGPPALEPPHNLVPVQLTSLRPDITQPPSNGT
 POEV1 GSTLTIRLR IETGTEPPVAMGPKVLAEQGPPALEPPHNLVPVQLTslrpditqppsnst

PERVA SGSVPT-----EPNITIKTGAKLFSLIQGAFOALNSTTPEATSSCWLC LASGPPYYEGMA
 POEVMSL SGSDPT-----ESNSTTKMGAKLFSLIQGAFOALNSTTPEATSSCWLC LASGPPYYEGMA
 TSUKUBA SGSDPT-----ESSSTTKMGAKLFSLIQGAFOALNSTTPEATSSCWLC LALGPPYYEGMA
 PERVB TGLIPTNTPRNSPGVPVKTGQRLFSLIQGAFOAINSTD PDATSSCWLC LSSGPPYYEGMA
 POEV1 TGLIPTNTPRNSPGVPVKTGQRLFSLIQGAFOAINSTD PDATSSCWLC LSSGPPYYEGMA

FIGURE 3



ATTORNEY DOCKET NO. 9013.43

DECLARATION, POWER OF ATTORNEY AND PETITION

As the below named inventor, I hereby declare:

my residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled _____

RETROVIRUS ASSAY _____, the specification of which
☐ is attached hereto.

☒ was filed on Feb. 15, 2002,

as Application Serial No. 10/069,082

and was amended on _____ (if applicable).

☒ was described and claimed in PCT International Application No. PCT/GB00/03159
and as amended under PCT Article 19 on _____ (if any).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the United States Patent & Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

PRIORITY CLAIM UNDER 35 USC § 119(a)-(d)

I hereby claim foreign priority benefits under Title 35, United States Code, §119(a)-(d) or §365(b) of any foreign applications(s) for patent or inventor's certificate, or §365(a) of any PCT International Applications designating at least one country other than the U.S. listed below and have also identified below any foreign application for patent or inventor's certificate or of any PCT International Applications designating at least one country other than the U.S. having a filing date before that of the application on which priority is claimed:

☐ no such applications have been filed

☒ application(s) listed below:

PRIOR FOREIGN APPLICATION(S)
Filed Within Twelve Months (Six Months For Design) Of This Application

			PRIORITY CLAIMED	
			YES	NO
9919604.0	United Kingdom	18 August 1999	<input checked="" type="checkbox"/>	<input type="checkbox"/>
(Number)	(Country)	(Day/month/year filed)		
<hr/>			<input type="checkbox"/>	<input type="checkbox"/>
(Number)	(Country)	(Day/month/year filed)		
<hr/>			<input type="checkbox"/>	<input type="checkbox"/>
(Number)	(Country)	(Day/month/year filed)		

CLAIM FOR BENEFIT OF PROVISIONAL APPLICATION UNDER 35 USC §119(e)

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States Provisional application listed below:

PROVISIONAL APPLICATION NO.

FILING DATE

CLAIM FOR BENEFIT OF EARLIER APPLICATIONS UNDER 35 USC §120

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s), or §365(c) of any PCT International Application(s) designating the U.S. listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose to the U.S. Patent & Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

(App. Serial No.)	(Filing date)	(Status) (patented, pending, abandoned)
(App. Serial No.)	(Filing date)	(Status) (patented, pending, abandoned)

PRIOR FOREIGN APPLICATIONS(Filed More Than Twelve Months (Six Months for Design) Prior To This Application)

_____ (Number)	_____ (Country)	_____ (Day/month/year filed)
-------------------	--------------------	---------------------------------

_____ (Number)	_____ (Country)	_____ (Day/month/year filed)
-------------------	--------------------	---------------------------------

_____ (Number)	_____ (Country)	_____ (Day/month/year filed)
-------------------	--------------------	---------------------------------

POWER OF ATTORNEY

I and I hereby appoint the practitioners associated with the Customer Number provided below as my attorneys, to prosecute this application and to transact all business in the United States Patent and Trademark Office connected therewith, and direct that all correspondence be addressed to that Customer Number:

Customer Number 20792

Send correspondence to: Kenneth D Sibley
Myers Bigel Sibley & Sajovec
Post Office Box 37428
Raleigh, NC 27627

Direct telephone calls to: Kenneth D Sibley
(919) 854 1400

Facsimile: (919) 854 1401

DECLARATION

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor Daniel GALBRAITH

100
Inventor's signature *D Galbraith*

Date *3/10/02* Citizenship British

Residence Glasgow, United Kingdom *GBX*

Post Office Address 513 Alexandra Parade, Glasgow, G31 3EW, United Kingdom

Full name of second, joint inventor, if any Helena KELLY

200
Inventor's signature *Helena T. Kelly*

Date *4th October 2002* Citizenship British

Residence Glasgow, United Kingdom *GBX*

Post Office Address Flat 1/2, 19 Kelvindale Gardens, Glasgow, G20 8DW, United Kingdom

Full name of third, joint inventor, if any Kenneth SMITH

300
Inventor's signature *Kenneth T Smith*

Date *4th October 2002* Citizenship British

Residence Glasgow, United Kingdom *GBX*

Post Office Address 5 Capelrig Lane, Newton Meams, Glasgow, G77 6XZ, United Kingdom



Rec'd PCT/PTO 30-OCT-2002
10/069082

SEQUENCE LISTING

Quip Technology Limited

<120> Retrovirus Assay

<130> P10890US

<140> US 10/069,082

<141> 2000-08-16

<150> GB 9919604.0

<151> 1999-08-18

<150> PCT/GB00/03159

<151> 2000-08-16

<160> 19

<170> PatentIn version 3.1

<210> 1

<211> 524

<212> PRT

<213> porcine endogenous retrovirus

<400> 1

Met Gly Gln Thr Val Thr Thr Pro Leu Ser Leu Thr Leu Asp His Trp
1 5 10 15

Thr Glu Val Arg Ser Arg Ala His Asn Leu Ser Val Gln Val Lys Lys
20 25 30

Gly Pro Trp Gln Thr Phe Cys Ala Ser Glu Trp Pro Thr Phe Asp Val
35 40 45

Gly Trp Pro Ser Glu Gly Thr Phe Asn Ser Glu Ile Ile Leu Ala Val
50 55 60

Lys Ala Ile Ile Phe Gln Thr Gly Pro Gly Ser His Pro Asp Gln Glu
65 70 75 80

Pro Tyr Ile Leu Thr Trp Gln Asp Leu Ala Glu Asp Pro Pro Pro Trp
85 90 95

Val Lys Pro Trp Leu Asn Lys Pro Arg Lys Pro Gly Pro Arg Ile Leu
100 105 110

Ala Leu Gly Glu Lys Asn Lys His Ser Ala Glu Lys Val Glu Pro Ser
115 120 125

Ser Ser Tyr Leu Pro Arg Asp Arg Gly Ala Ala Asp Leu Ala Gly Thr
130 135 140

Pro Thr Cys Ser Pro Thr Pro Leu Ser Ser Thr Gly Cys Cys Glu Gly
145 150 155 160

Thr Ser Ala Pro Pro Gly Ala Pro Val Val Glu Gly Pro Ala Ala Gly
165 170 175

Thr Arg Ser Arg Arg Gly Ala Thr Pro Glu Arg Thr Asp Glu Ile Ala
180 185 190

Ile Leu Pro Leu Arg Thr Tyr Gly Pro Pro Met Pro Gly Gly Gln Leu
195 200 205

Gln Pro Leu Gln Tyr Trp Pro Phe Ser Ser Ala Asp Leu Tyr Asn Trp
210 215 220

Lys Thr Asn His Pro Pro Phe Ser Glu Asp Pro Gln Arg Leu Thr Gly
225 230 235 240

Leu Val Glu Ser Leu Met Phe Ser His Gln Pro Thr Trp Asp Asp Cys
245 250 255

Gln Gln Leu Leu Gln Thr Leu Phe Thr Thr Glu Glu Arg Glu Arg Ile
260 265 270

Leu Leu Glu Ala Arg Lys Asn Val Pro Gly Ala Asp Gly Arg Pro Thr
275 280 285

Gln Leu Gln Asn Glu Ile Asp Met Gly Phe Pro Leu Thr Arg Pro Gly
290 295 300

Trp Asp Tyr Asn Thr Ala Glu Gly Arg Glu Ser Leu Lys Ile Tyr Arg
305 310 315 320

Gln Ala Leu Val Ala Gly Leu Arg Gly Ala Ser Arg Arg Pro Thr Asn
325 330 335

Leu Ala Lys Val Arg Glu Val Met Gln Gly Pro Asn Glu Pro Pro Ser
340 345 350

Val Phe Leu Glu Arg Leu Met Glu Ala Phe Arg Arg Phe Thr Pro Phe
355 360

Asp Pro Thr Ser Glu Ala Gln Lys Ala Ser Val Ala Leu Ala Phe Ile
370 375 380

Gly Gln Ser Ala Leu Asp Ile Arg Lys Lys Leu Gln Arg Leu Glu Gly
385 390 395 400

Leu Gln Glu Ala Glu Leu Arg Asp Leu Val Arg Glu Ala Glu Lys Val
405 410 415

Tyr Tyr Arg Arg Glu Thr Glu Glu Glu Lys Glu Gln Arg Lys Glu Lys
420 425 430

Glu Arg Glu Glu Arg Glu Glu Arg Arg Asp Arg Arg Gln Glu Lys Asn
435 440 445

Leu Thr Lys Ile Leu Ala Ala Val Val Glu Gly Lys Ser Ser Arg Glu
450 455 460

Arg Glu Arg Asp Phe Arg Lys Ile Arg Ser Gly Pro Arg Gln Ser Gly
465 470 475 480

Asn Leu Gly Asn Arg Thr Pro Leu Asp Lys Asp Gln Cys Ala Tyr Cys
485 490 495

Lys Glu Lys Gly His Trp Ala Arg Asn Cys Pro Lys Lys Gly Asn Lys
500 505 510

Gly Pro Lys Val Leu Ala Leu Glu Glu Asp Lys Asp
515 520

<210> 2

<211> 660

<212> PRT

<213> porcine endogenous retrovirus Type PERV A

<400> 2

Met His Pro Thr Leu Ser Arg Arg His Leu Pro Ile Arg Gly Gly Lys
1 5 10 15

Pro Lys Arg Leu Lys Ile Pro Leu Ser Phe Ala Ser Ile Ala Trp Phe
20 25 30

Leu Thr Leu Ser Ile Thr Pro Gln Val Asn Gly Lys Arg Leu Val Asp
35 40 45

Ser Pro Asn Ser His Lys Pro Leu Ser Leu Thr Trp Leu Leu Thr Asp
 50 55 60
 Ser Gly Thr Gly Ile Asn Ile Asn Ser Thr Gln Gly Glu Ala Pro Leu
 65 70 75 80
 Gly Thr Trp Trp Pro Glu Leu Tyr Val Cys Leu Arg Ser Val Ile Pro
 85 90 95
 Gly Leu Asn Asp Gln Ala Thr Pro Pro Asp Val Leu Arg Ala Tyr Gly
 100 105 110
 Phe Tyr Val Cys Pro Gly Pro Pro Asn Asn Glu Glu Tyr Cys Gly Asn
 115 120 125
 Pro Gln Asp Phe Phe Cys Lys Gln Trp Ser Cys Ile Thr Ser Asn Asp
 130 135 140
 Gly Asn Trp Lys Trp Pro Val Ser Gln Gln Asp Arg Val Ser Tyr Ser
 145 150 155 160
 Phe Val Asn Asn Pro Thr Ser Tyr Asn Gln Phe Asn Tyr Gly His Gly
 165 170 175
 Arg Trp Lys Asp Trp Gln Gln Arg Val Gln Lys Asp Val Arg Asn Lys
 180 185 190
 Gln Ile Ser Cys His Ser Leu Asp Leu Asp Tyr Leu Lys Ile Ser Phe
 195 200 205
 Thr Glu Lys Gly Lys Gln Glu Asn Ile Gln Lys Trp Val Asn Gly Ile
 210 215 220
 Ser Trp Gly Ile Val Tyr Tyr Gly Gly Ser Gly Arg Lys Lys Gly Ser
 225 230 235 240
 Val Leu Thr Ile Arg Leu Arg Ile Glu Thr Gln Met Glu Pro Pro Val
 245 250 255
 Ala Ile Gly Pro Asn Lys Gly Leu Ala Glu Gln Gly Pro Pro Ile Gln
 260 265 270
 Glu Gln Arg Pro Ser Pro Asn Pro Ser Asp Tyr Asn Thr Thr Ser Gly
 275 280 285
 Ser Val Pro Thr Glu Pro Asn Ile Thr Ile Lys Thr Gly Ala Lys Leu
 290 295 300
 Phe Ser Leu Ile Gln Gly Ala Phe Gln Ala Leu Asn Ser Thr Thr Pro
 305 310 315 320
 Glu Ala Thr Ser Ser Cys Trp Leu Cys Leu Ala Ser Gly Pro Pro Tyr
 325 330 335

Tyr	Glu	Gly	Met	Ala	Arg	Gly	Gly	Lys	Phe	Asn	Val	Thr	Lys	Glu	His
			340					345					350		
Arg	Asp	Gln	Cys	Thr	Trp	Gly	Ser	Gln	Asn	Lys	Leu	Thr	Leu	Thr	Glu
		355					360					365			
Val	Ser	Gly	Lys	Gly	Thr	Cys	Ile	Gly	Met	Val	Pro	Pro	Ser	His	Gln
		370				375					380				
His	Leu	Cys	Asn	His	Thr	Glu	Ala	Phe	Asn	Arg	Thr	Ser	Glu	Ser	Gln
		385			390					395					400
Tyr	Leu	Val	Pro	Gly	Tyr	Asp	Arg	Trp	Trp	Ala	Cys	Asn	Thr	Gly	Leu
				405					410					415	
Thr	Pro	Cys	Val	Ser	Thr	Leu	Val	Phe	Asn	Gln	Thr	Lys	Asp	Phe	Cys
			420					425					430		
Val	Met	Val	Gln	Ile	Val	Pro	Arg	Val	Tyr	Tyr	Tyr	Pro	Glu	Lys	Ala
		435					440					445			
Val	Leu	Asp	Glu	Tyr	Asp	Tyr	Arg	Tyr	Asn	Arg	Pro	Lys	Arg	Glu	Pro
		450				455					460				
Ile	Ser	Leu	Thr	Leu	Ala	Val	Met	Leu	Gly	Leu	Gly	Val	Ala	Ala	Gly
		465			470					475					480
Val	Gly	Thr	Gly	Thr	Ala	Ala	Leu	Ile	Thr	Gly	Pro	Gln	Gln	Leu	Glu
				485					490					495	
Lys	Gly	Leu	Ser	Asn	Leu	His	Arg	Ile	Val	Thr	Glu	Asp	Leu	Gln	Ala
			500					505					510		
Leu	Glu	Lys	Ser	Val	Ser	Asn	Leu	Glu	Glu	Ser	Leu	Thr	Ser	Leu	Ser
		515					520					525			
Glu	Val	Val	Leu	Gln	Asn	Arg	Arg	Gly	Leu	Asp	Leu	Leu	Phe	Leu	Lys
		530				535					540				
Glu	Gly	Gly	Leu	Cys	Val	Ala	Leu	Lys	Glu	Glu	Cys	Cys	Phe	Tyr	Val
		545			550					555					560
Asp	His	Ser	Gly	Ala	Ile	Arg	Asp	Ser	Met	Ser	Lys	Leu	Arg	Glu	Arg
				565					570					575	
Leu	Glu	Arg	Arg	Arg	Arg	Glu	Arg	Glu	Ala	Asp	Gln	Gly	Trp	Phe	Glu
			580						585				590		
Gly	Trp	Phe	Asn	Arg	Ser	Pro	Trp	Met	Thr	Thr	Leu	Leu	Ser	Ala	Leu
		595					600					605			

Thr Gly Pro Leu Val Val Leu Leu Leu Leu Thr Val Gly Pro Cys
610 615 620

Leu Ile Asn Arg Phe Val Ala Phe Val Arg Glu Arg Val Ser Ala Val
625 630 635 640

Gln Ile Met Val Leu Arg Gln Gln Tyr Gln Gly Leu Leu Ser Gln Gly
645 650 655

Glu Thr Asp Leu
660

<210> 3

<211> 638

<212> PRT

<213> porcine endogenous retrovirus Type PERV MSL

<400> 3

Met His Pro Thr Leu Asn Arg Arg His Leu Pro Ile Arg Gly Gly Lys
1 5 10 15

Pro Lys Arg Leu Lys Ile Pro Leu Ser Phe Ala Ser Ile Ala Trp Phe
20 25 30

Leu Thr Leu Ser Ile Thr Ser Gln Thr Asn Gly Met Arg Ile Gly Asp
35 40 45

Ser Leu Asn Ser His Lys Pro Leu Ser Leu Thr Trp Leu Ile Thr Asp
50 55 60

Ser Gly Thr Gly Ile Asn Ile Asn Asn Thr Gln Gly Glu Ala Pro Leu
65 70 75 80

Gly Thr Trp Trp Pro Asp Leu Tyr Val Cys Leu Arg Ser Val Ile Pro
85 90 95

Ser Leu Thr Ser Pro Pro Asp Ile Leu His Ala His Gly Phe Tyr Val
100 105 110

Cys Pro Gly Pro Pro Asn Asn Gly Lys His Cys Gly Asn Pro Arg Asp
115 120 125

Phe Phe Cys Lys Gln Trp Asn Cys Val Thr Ser Asn Asp Gly Tyr Trp
130 135 140

Lys Trp Pro Thr Ser Gln Gln Asp Arg Val Ser Phe Ser Tyr Val Asn
145 150 155 160

Thr Tyr Thr Ser Ser Gly Gln Phe Asn Tyr Leu Thr Trp Ile Arg Thr
165 170 175

Gly Ser Pro Lys Cys Ser Ser Asp Leu Asp Tyr Leu Lys Ile Ser
180 185

Phe Thr Glu Lys Gly Lys Gln Glu Asn Ile Leu Lys Trp Val Asn Gly
195 200

Met Ser Trp Gly Met Val Tyr Tyr Gly Gly Ser Gly Lys Gln Pro Gly
210 215 220

Ser Ile Leu Thr Ile Arg Leu Lys Ile Asn Gln Leu Glu Pro Pro Met
225 230 235 240

Ala Ile Gly Pro Asn Thr Val Leu Thr Gly Gln Arg Pro Pro Thr Gln
245 250 255

Gly Pro Gly Pro Ser Ser Asn Ile Thr Ser Gly Ser Asp Pro Thr Glu
260 265 270

Ser Asn Ser Thr Thr Lys Met Gly Ala Lys Leu Phe Ser Leu Ile Gln
275 280 285

Gly Ala Phe Gln Ala Leu Asn Ser Thr Thr Pro Glu Ala Thr Ser Ser
290 295 300

Cys Trp Leu Cys Leu Ala Ser Gly Pro Pro Tyr Tyr Glu Gly Met Ala
305 310 315 320

Arg Arg Gly Lys Phe Asn Val Thr Lys Glu His Arg Asp Gln Cys Thr
325 330 335

Trp Gly Ser Gln Asn Lys Leu Thr Leu Thr Glu Val Ser Gly Lys Gly
340 345 350

Thr Cys Ile Gly Lys Val Pro Pro Ser His Gln His Leu Cys Asn His
355 360 365

Thr Glu Ala Phe Asn Gln Thr Ser Glu Ser Gln Tyr Leu Val Pro Gly
370 375 380

Tyr Asp Arg Trp Trp Ala Cys Asn Thr Gly Leu Thr Pro Cys Val Ser
385 390 395 400

Thr Leu Val Phe Asn Gln Thr Lys Asp Phe Cys Ile Met Val Gln Ile
405 410 415

Val Pro Arg Val Tyr Tyr Tyr Pro Glu Lys Ala Ile Leu Asp Glu Tyr
420 425 430

Asp Tyr Arg Asn His Arg Gln Lys Arg Glu Pro Ile Ser Leu Thr Leu
435 440 445

Ala Val Met Leu Gly Leu Gly Val Ala Ala Gly Val Gly Thr Gly Thr
450 455 460

Ala Ala Leu Val Thr Gly Pro Gln Gln Leu Glu Thr Gly Leu Ser Asn
465 470 475 480

Leu His Arg Ile Val Thr Glu Asp Leu Gln Ala Leu Glu Lys Ser Val
485 490 495

Ser Asn Leu Glu Glu Ser Leu Thr Ser Leu Ser Glu Val Val Leu Gln
500 505 510

Asn Arg Arg Gly Leu Asp Leu Leu Phe Leu Lys Glu Gly Gly Leu Cys
515 520 525

Val Ala Leu Lys Glu Glu Cys Cys Phe Tyr Val Asp His Ser Gly Ala
530 535 540

Ile Arg Asp Ser Met Asn Lys Leu Arg Glu Arg Leu Glu Lys Arg Arg
545 550 555 560

Arg Glu Lys Glu Thr Thr Gln Gly Trp Phe Glu Gly Trp Phe Asn Arg
565 570 575

Ser Leu Trp Leu Ala Thr Leu Leu Ser Ala Leu Thr Gly Pro Leu Ile
580 585 590

Val Leu Leu Leu Leu Thr Val Gly Pro Cys Ile Ile Asn Lys Leu
595 600 605

Ile Ala Phe Ile Arg Glu Arg Ile Ser Ala Val Gln Ile Met Val Leu
610 615 620

Arg Gln Gln Tyr Gln Ser Pro Ser Ser Arg Glu Ala Gly Arg
625 630 635

<210> 4

<211> 638

<212> PRT

<213> porcine endogenous retrovirus Type Tsukuba

<400> 4

Met His Pro Thr Leu Ser Arg Arg His Leu Pro Ile Arg Gly Gly Lys
1 5 10 15

Pro Lys Arg Leu Lys Ile Pro Leu Ser Phe Ala Ser Ile Ala Trp Phe
20 25 30

Leu Thr Leu Ser Ile Thr Ser Gln Thr Asn Gly Met Arg Ile Gly Asp
35 40 45

Ser Leu Asn Ser His Lys Pro Leu Ser Leu Thr Trp Leu Ile Thr Asp
50 55 60

Ser Gly Thr Gly Ile Asn Ile Asn Asn Thr Gln Gly Glu Ala Pro Leu
65 70 75 80

Gly Thr Trp Trp Pro Asp Leu Tyr Val Cys Leu Arg Ser Val Ile Pro
85 90 95

Ser Leu Thr Ser Pro Pro Asp Ile Leu His Ala His Gly Phe Tyr Val
100 105 110

Cys Pro Gly Pro Pro Asn Asn Gly Lys His Cys Gly Asn Pro Arg Asp
115 120 125

Phe Phe Cys Lys Gln Trp Asn Cys Val Thr Ser Asn Asp Gly Tyr Trp
130 135 140

Lys Trp Pro Thr Ser Gln Gln Asp Arg Val Ser Phe Ser Tyr Val Asn
145 150 155 160

Thr Tyr Thr Ser Ser Gly Gln Phe Asn Tyr Leu Thr Trp Ile Arg Thr
165 170 175

Gly Ser Pro Lys Cys Ser Pro Ser Asp Leu Asp Tyr Leu Lys Ile Ser
180 185 190

Phe Thr Glu Lys Gly Lys Gln Glu Asn Ile Leu Lys Trp Val Asn Gly
195 200 205

Met Ser Trp Gly Met Val Tyr Tyr Gly Gly Ser Gly Lys Gln Pro Gly
210 215 220

Ser Ile Leu Thr Ile Arg Leu Lys Ile Asn Gln Leu Glu Pro Pro Met
225 230 235 240

Ala Ile Gly Pro Asn Thr Val Leu Thr Gly Gln Arg Pro Pro Thr Gln
245 250 255

Gly Pro Gly Pro Ser Ser Asn Ile Thr Ser Gly Ser Asp Pro Thr Glu
260 265 270

Ser Ser Ser Thr Thr Lys Met Gly Ala Lys Leu Phe Ser Leu Ile Gln
275 280 285

Gly Ala Phe Gln Ala Leu Asn Ser Thr Thr Pro Glu Ala Thr Ser Ser
290 295 300

Cys Trp Leu Cys Leu Ala Leu Gly Pro Pro Tyr Tyr Glu Gly Met Ala
305 310 315 320

Arg Arg Gly Lys Phe Asn Val Thr Lys Glu His Arg Asp Gln Cys Thr
325 330

Trp Gly Ser Gln Asn Lys Leu Thr Leu Thr Glu Val Ser Gly Lys Gly
340 345 350

Thr Cys Ile Gly Lys Val Pro Pro Ser His Gln His Leu Cys Asn His
355 360 365

Thr Glu Ala Phe Asn Gln Thr Ser Glu Ser Gln Tyr Leu Val Pro Gly
370 375 380

Tyr Asp Arg Trp Trp Ala Cys Asn Thr Gly Leu Thr Pro Cys Val Ser
385 390 395 400

Thr Leu Val Phe Asn Gln Thr Lys Asp Phe Cys Ile Met Val Gln Ile
405 410 415

Val Pro Arg Val Tyr Tyr Tyr Pro Glu Lys Ala Ile Leu Asp Glu Tyr
420 425 430

Asp Tyr Arg Asn His Arg Gln Lys Arg Glu Pro Ile Ser Leu Thr Leu
435 440 445

Ala Val Met Leu Gly Leu Gly Val Ala Ala Gly Val Gly Thr Gly Thr
450 455 460

Ala Ala Leu Val Thr Gly Pro Gln Gln Leu Glu Thr Gly Leu Ser Asn
465 470 475 480

Leu His Arg Ile Val Thr Glu Asp Leu Gln Ala Leu Glu Lys Ser Val
485 490 495

Ser Asn Leu Glu Glu Ser Leu Thr Ser Leu Ser Glu Val Val Leu Gln
500 505 510

Asn Arg Arg Gly Leu Asp Leu Leu Phe Leu Lys Glu Gly Gly Leu Cys
515 520 525

Val Ala Leu Lys Glu Glu Cys Cys Phe Tyr Val Asp His Ser Gly Ala
530 535 540

Ile Arg Asp Ser Met Asn Lys Leu Arg Glu Arg Leu Glu Lys Arg Arg
545 550 555 560

Arg Glu Lys Glu Thr Thr Gln Gly Trp Phe Glu Gly Trp Phe Asn Arg
565 570 575

Ser Pro Trp Leu Ala Thr Leu Leu Ser Ala Leu Thr Gly Pro Leu Ile
580 585 590

Val Leu Leu Leu Leu Thr Val Gly Pro Cys Ile Ile Asn Lys Leu
595 600 605

Ile Ala Phe Ile Arg Glu Arg Ile Ser Ala Val Gln Ile Met Val Leu
610 615 620

Arg Gln Gln Tyr Gln Ser Pro Ser Ser Arg Glu Ala Gly Arg
625 630 635

<210> 5

<211> 657

<212> PRT

<213> porcine endogenous retrovirus Type PERV B

<400> 5

Met His Pro Thr Leu Ser Trp Arg His Leu Pro Thr Arg Gly Gly Glu
1 5 10 15

Pro Lys Arg Leu Arg Ile Pro Leu Ser Phe Ala Ser Ile Ala Trp Phe
20 25 30

Leu Thr Leu Thr Ile Thr Pro Gln Ala Ser Ser Lys Arg Leu Ile Asp
35 40 45

Ser Ser Asn Pro His Arg Pro Leu Ser Leu Thr Trp Leu Ile Ile Asp
50 55 60

Pro Asp Thr Gly Val Thr Val Asn Ser Thr Arg Gly Val Ala Pro Arg
65 70 75 80

Gly Thr Trp Trp Pro Glu Leu His Phe Cys Leu Arg Leu Ile Asn Pro
85 90 95

Ala Val Lys Ser Thr Pro Pro Asn Leu Val Arg Ser Tyr Gly Phe Tyr
100 105 110

Cys Cys Pro Gly Thr Glu Lys Glu Lys Tyr Cys Gly Gly Ser Gly Glu
115 120 125

Ser Phe Cys Arg Arg Trp Ser Cys Val Thr Ser Asn Asp Gly Asp Trp
130 135 140

Lys Trp Pro Ile Ser Leu Gln Asp Arg Val Lys Phe Ser Phe Val Asn
145 150 155 160

Ser Gly Pro Gly Lys Tyr Lys Val Met Lys Leu Tyr Lys Asp Lys Ser
165 170 175

Cys Ser Pro Ser Asp Leu Asp Tyr Leu Lys Ile Ser Phe Thr Glu Lys
180 185 190

Gly Lys Gln Glu Asn Ile Gln Lys Trp Ile Asn Gly Met Ser Trp Gly
 195 200
 Ile Val Phe Tyr Lys Tyr Gly Gly Gly Ala Gly Ser Thr Leu Thr Ile
 210 215 220
 Arg Leu Arg Ile Glu Thr Gly Thr Glu Pro Pro Val Ala Val Gly Pro
 225 230 235 240
 Asp Lys Val Leu Ala Glu Gln Gly Pro Pro Ala Leu Glu Pro Pro His
 245 250 255
 Asn Leu Pro Val Pro Gln Leu Thr Ser Leu Arg Pro Asp Ile Thr Gln
 260 265 270
 Pro Pro Ser Asn Gly Thr Thr Gly Leu Ile Pro Thr Asn Thr Pro Arg
 275 280 285
 Asn Ser Pro Gly Val Pro Val Lys Thr Gly Gln Arg Leu Phe Ser Leu
 290 295 300
 Ile Gln Gly Ala Phe Gln Ala Ile Asn Ser Thr Asp Pro Asp Ala Thr
 305 310 315 320
 Ser Ser Cys Trp Leu Cys Leu Ser Ser Gly Pro Pro Tyr Tyr Glu Gly
 325 330 335
 Met Ala Lys Glu Gly Lys Phe Asn Val Thr Lys Glu His Arg Asn Gln
 340 345 350
 Cys Thr Trp Gly Ser Arg Asn Lys Leu Thr Leu Thr Glu Val Ser Gly
 355 360 365
 Lys Gly Thr Cys Ile Gly Lys Ala Pro Pro Ser His Gln His Leu Cys
 370 375 380
 Tyr Ser Thr Val Val Tyr Glu Gln Ala Ser Glu Asn Gln Tyr Leu Val
 385 390 395 400
 Pro Gly Tyr Asn Arg Trp Trp Ala Cys Asn Thr Gly Leu Thr Pro Cys
 405 410 415
 Val Ser Thr Ser Val Phe Asn Gln Ser Lys Asp Phe Cys Val Met Val
 420 425 430
 Gln Ile Val Pro Arg Val Tyr Tyr His Pro Glu Glu Val Val Leu Asp
 435 440 445
 Glu Tyr Asp Tyr Arg Tyr Asn Arg Pro Lys Arg Glu Pro Val Ser Leu
 450 455 460
 Thr Leu Ala Val Met Leu Gly Leu Gly Thr Ala Val Gly Val Gly Thr
 465 470 475 480

Gly Thr Ala Ala Leu Ile Thr Gly Pro Gln Gln Leu Glu Lys Gly Leu
485 490 495

Gly Glu Leu His Ala Ala Met Thr Glu Asp Leu Arg Ala Leu Glu Glu
500 505 510

Ser Val Ser Asn Leu Glu Glu Ser Leu Thr Ser Leu Ser Glu Val Val
515 520 525

Leu Gln Asn Arg Arg Gly Leu Asp Leu Leu Phe Leu Arg Glu Gly Gly
530 535 540

Leu Cys Ala Ala Leu Lys Glu Glu Cys Cys Phe Tyr Val Asp His Ser
545 550 555 560

Gly Ala Ile Arg Asp Ser Met Ser Lys Leu Arg Glu Arg Leu Glu Arg
565 570 575

Arg Arg Arg Glu Arg Glu Ala Asp Gln Gly Trp Phe Glu Gly Trp Phe
580 585 590

Asn Arg Ser Pro Trp Met Thr Thr Leu Leu Ser Ala Leu Thr Gly Pro
595 600 605

Leu Val Val Leu Leu Leu Leu Leu Thr Val Gly Pro Cys Leu Ile Asn
610 615 620

Arg Phe Val Ala Phe Val Arg Glu Arg Val Ser Ala Val Gln Ile Met
625 630 635 640

Val Leu Arg Gln Gln Tyr Gln Gly Leu Leu Ser Gln Gly Glu Thr Asp
645 650 655

Leu

<210> 6

<211> 656

<212> PRT

<213> porcine endogenous retrovirus Type PoEV1

<400> 6

Met His Pro Thr Leu Ser Arg Arg His Leu Pro Thr Arg Gly Gly Glu
1 5 10 15

Pro Lys Arg Leu Arg Ile Pro Leu Ser Phe Ala Ser Ile Ala Trp Phe
20 25 30

Leu Thr Leu Thr Ile Thr Pro Gln Ala Ser Ser Lys Arg Leu Ile Asp
 35 40
 Ser Ser Asn Pro His Arg Pro Leu Ser Leu Thr Trp Leu Ile Ile Asp
 50 55 60
 Pro Asp Thr Gly Val Thr Val Asn Ser Thr Arg Gly Val Ala Pro Arg
 65 70 75 80
 Gly Thr Trp Trp Pro Glu Leu His Phe Cys Leu Arg Leu Ile Asn Pro
 85 90 95
 Ala Val Lys Ser Thr Pro Pro Asn Leu Val Arg Ser Tyr Gly Phe Tyr
 100 105 110
 Cys Cys Pro Gly Thr Glu Lys Glu Lys Tyr Cys Gly Gly Ser Gly Glu
 115 120 125
 Ser Phe Cys Arg Arg Trp Ser Cys Val Thr Ser Asn Asp Gly Asp Trp
 130 135 140
 Lys Trp Pro Ile Ser Leu Gln Asp Arg Val Lys Phe Ser Phe Val Asn
 145 150 155 160
 Ser Gly Pro Gly Lys Tyr Lys Met Met Lys Leu Tyr Lys Asp Lys Ser
 165 170 175
 Cys Ser Pro Ser Asp Leu Asp Tyr Leu Lys Ile Ser Phe Thr Glu Arg
 180 185 190
 Lys Thr Gly Lys Tyr Ser Lys Val Asp Lys Trp Tyr Glu Leu Gly Asn
 195 200 205
 Ser Phe Leu Leu Tyr Gly Gly Gly Ala Gly Ser Thr Leu Thr Ile Arg
 210 215 220
 Leu Arg Ile Glu Thr Gly Thr Glu Pro Pro Val Ala Met Gly Pro Asp
 225 230 235 240
 Lys Val Leu Ala Glu Gln Gly Pro Pro Ala Leu Glu Pro Pro His Asn
 245 250 255
 Leu Pro Val Pro Gln Leu Thr Ser Leu Arg Pro Asp Ile Thr Gln Pro
 260 265 270
 Pro Ser Asn Ser Thr Thr Gly Leu Ile Pro Thr Asn Thr Pro Arg Asn
 275 280 285
 Ser Pro Gly Val Pro Val Lys Thr Gly Gln Arg Leu Phe Ser Leu Ile
 290 295 300
 Gln Gly Ala Phe Gln Ala Ile Asn Ser Thr Asp Pro Asp Ala Thr Ser
 305 310 315 320

Ser Cys Trp Leu Cys₃₂₅ Leu Ser Ser Gly₃₃₀ Pro Pro Tyr Tyr Glu₃₃₅ Met

Ala Lys Glu Arg₃₄₀ Lys Phe Asn Val Thr₃₄₅ Lys Glu His Arg Asn₃₅₀ Gln Cys

Thr Trp Gly₃₅₅ Ser Arg Asn Lys Leu₃₆₀ Thr Leu Thr Glu₃₆₅ Val Ser Gly Lys

Gly Thr Cys Ile Gly Lys Ala₃₇₅ Pro Pro Ser His Gln₃₈₀ His Leu Cys Tyr

Ser Thr Val Val Tyr Glu₃₉₀ Gln Ala Ser Glu₃₉₅ Asn Gln Tyr Leu Val Pro₄₀₀

Gly Tyr Asn Arg Trp₄₀₅ Trp Ala Cys Asn Thr₄₁₀ Gly Leu Thr Pro Cys₄₁₅ Val

Ser Thr Ser Val₄₂₀ Phe Asn Gln Ser Lys₄₂₅ Asp Phe Cys Val Met₄₃₀ Val Gln

Ile Val Pro Arg Val Tyr Tyr His₄₄₀ Pro Glu Glu Val₄₄₅ Val Leu Asp Glu

Tyr Asp Tyr Arg Tyr Asn Arg₄₅₅ Pro Lys Arg Glu₄₆₀ Pro Val Ser Leu Thr

Leu Ala Val Met Leu Gly₄₇₀ Leu Gly Thr Ala Val₄₇₅ Gly Val Gly Thr Gly₄₈₀

Thr Ala Ala Leu Ile₄₈₅ Thr Gly Pro Gln Gln₄₉₀ Leu Glu Lys Gly₄₉₅ Leu Gly

Glu Leu His Ala₅₀₀ Ala Met Thr Glu Asp₅₀₅ Leu Arg Ala Leu Lys₅₁₀ Glu Ser

Val Ser Asn₅₁₅ Leu Glu Glu Ser Leu Thr Ser Leu Ser Glu₅₂₀ Val Val Leu

Gln Asn Arg Arg Gly Leu Asp₅₃₅ Leu Leu Phe Leu Arg₅₄₀ Glu Gly Gly Leu

Cys Ala Ala Leu Lys Glu₅₅₀ Glu Cys Cys Phe Tyr₅₅₅ Val Asp His Ser Gly₅₆₀

Ala Ile Arg Asp Ser₅₆₅ Met Asn Lys Leu Arg₅₇₀ Lys Lys Leu Glu Arg₅₇₅ Glu

Arg Arg Glu Arg₅₈₀ Glu Ala Asp Gln Gly₅₈₅ Trp Phe Glu Gly Trp₅₉₀ Phe Asn

Arg Ser Pro Trp Met Thr Thr Leu Leu Ser Ala Leu Thr Gly Pro Leu
595 600 605

Val Val Leu Leu Leu Leu Leu Thr Val Gly Pro Cys Leu Ile Asn Arg
610 615 620

Phe Val Ala Phe Val Arg Glu Arg Val Ser Ala Val Gln Ile Met Val
625 630 635 640

Leu Arg Gln Gln Tyr Gln Gly Leu Leu Ser Gln Gly Glu Thr Asp Leu
645 650 655

<210> 7

<211> 221

<212> PRT

<213> porcine endogenous retrovirus Type PERV A

<400> 7

Asn Asn Glu Glu Tyr Cys Gly Asn Pro Gln Asp Phe Phe Cys Lys Gln
1 5 10 15

Trp Ser Cys Ile Thr Ser Asn Asp Gly Asn Trp Lys Trp Pro Val Ser
20 25 30

Gln Gln Asp Arg Val Ser Tyr Ser Phe Val Asn Asn Pro Thr Ser Tyr
35 40 45

Asn Gln Phe Asn Tyr Gly His Gly Arg Trp Lys Asp Trp Gln Gln Arg
50 55 60

Val Gln Lys Asp Val Arg Asn Lys Gln Ile Ser Cys His Ser Leu Asp
65 70 75 80

Leu Asp Tyr Leu Lys Ile Ser Phe Thr Glu Lys Gly Lys Gln Glu Asn
85 90 95

Ile Gln Lys Trp Val Asn Gly Ile Ser Trp Gly Ile Val Tyr Tyr Gly
100 105 110

Gly Ser Gly Arg Lys Lys Gly Ser Val Leu Thr Ile Arg Leu Arg Ile
115 120 125

Glu Thr Gln Met Glu Pro Pro Val Ala Ile Gly Pro Asn Lys Gly Leu
130 135 140

Ala Glu Gln Gly Pro Pro Ile Gln Glu Gln Arg Pro Ser Pro Asn Pro
145 150 155 160

Ser Asp Tyr Asn Thr Thr Ser Gly Ser Val Pro Thr Glu Pro Asn Ile
165 170 175

Thr Ile Lys Thr Gly Ala Lys Leu Phe Ser Leu Ile Gln Gly Ala Phe
180 185 190

Gln Ala Leu Asn Ser Thr Thr Pro Glu Ala Thr Ser Ser Cys Trp Leu
195 200 205

Cys Leu Ala Ser Gly Pro Pro Tyr Tyr Glu Gly Met Ala
210 215 220

<210> 8

<211> 203

<212> PRT

<213> porcine endogenous retrovirus Type PERV MSL

<400> 8

Asn Asn Gly Lys His Cys Gly Asn Pro Arg Asp Phe Phe Cys Lys Gln
1 5 10 15

Trp Asn Cys Val Thr Ser Asn Asp Gly Tyr Trp Lys Trp Pro Thr Ser
20 25 30

Gln Gln Asp Arg Val Ser Phe Ser Tyr Val Asn Thr Tyr Thr Ser Ser
35 40 45

Gly Gln Phe Asn Tyr Leu Thr Trp Ile Arg Thr Gly Ser Pro Lys Cys
50 55 60

Ser Pro Ser Asp Leu Asp Tyr Leu Lys Ile Ser Phe Thr Glu Lys Gly
65 70 75 80

Lys Gln Glu Asn Ile Leu Lys Trp Val Asn Gly Met Ser Trp Gly Met
85 90 95

Val Tyr Tyr Gly Gly Ser Gly Lys Gln Pro Gly Ser Ile Leu Thr Ile
100 105 110

Arg Leu Lys Ile Asn Gln Leu Glu Pro Pro Met Ala Ile Gly Pro Asn
115 120 125

Thr Val Leu Thr Gly Gln Arg Pro Pro Thr Gln Gly Pro Gly Pro Ser
130 135 140

Ser Asn Ile Thr Ser Gly Ser Asp Pro Thr Glu Ser Asn Ser Thr Thr
145 150 155 160

Lys Met Gly Ala Lys Leu Phe Ser Leu Ile Gln Gly Ala Phe Gln Ala
165 170 175

Leu Asn Ser Thr Thr Pro Glu Ala Thr Ser Ser Cys Trp Leu Cys Leu
180 185 190

Ala Ser Gly Pro Pro Tyr Tyr Glu Gly Met Ala
195 200

<210> 9

<211> 203

<212> PRT

<213> porcine endogenous retrovirus Type Tsukuba

<400> 9

Asn Asn Gly Lys His Cys Gly Asn Pro Arg Asp Phe Phe Cys Lys Gln
1 5 10 15

Trp Asn Cys Val Thr Ser Asn Asp Gly Tyr Trp Lys Trp Pro Thr Ser
20 25 30

Gln Gln Asp Arg Val Ser Phe Ser Tyr Val Asn Thr Tyr Thr Ser Ser
35 40 45

Gly Gln Phe Asn Tyr Leu Thr Trp Ile Arg Thr Gly Ser Pro Lys Cys
50 55 60

Ser Pro Ser Asp Leu Asp Tyr Leu Lys Ile Ser Phe Thr Glu Lys Gly
65 70 75 80

Lys Gln Glu Asn Ile Leu Lys Trp Val Asn Gly Met Ser Trp Gly Met
85 90 95

Val Tyr Tyr Gly Gly Ser Gly Lys Gln Pro Gly Ser Ile Leu Thr Ile
100 105 110

Arg Leu Lys Ile Asn Gln Leu Glu Pro Pro Met Ala Ile Gly Pro Asn
115 120 125

Thr Val Leu Thr Gly Gln Arg Pro Pro Thr Gln Gly Pro Gly Pro Ser
130 135 140

Ser Asn Ile Thr Ser Gly Ser Asp Pro Thr Glu Ser Ser Ser Thr Thr
145 150 155 160

Lys Met Gly Ala Lys Leu Phe Ser Leu Ile Gln Gly Ala Phe Gln Ala
165 170 175

Leu Asn Ser Thr Thr Pro Glu Ala Thr Ser Ser Cys Trp Leu Cys Leu
180 185 190

Ala Leu Gly Pro Pro Tyr Tyr Glu Gly Met Ala
195 200

<210> 10

<211> 221

<212> PRT

<213> porcine endogenous retrovirus Type PERV B

<400> 10

Glu Lys Glu Lys Tyr Cys Gly Gly Ser Gly Glu Ser Phe Cys Arg Arg
1 5 10

Trp Ser Cys Val Thr Ser Asn Asp Gly Asp Trp Lys Trp Pro Ile Ser
20 25 30

Leu Gln Asp Arg Val Lys Phe Ser Phe Val Asn Ser Gly Pro Gly Lys
35 40 45

Tyr Lys Val Met Lys Leu Tyr Lys Asp Lys Ser Cys Ser Pro Ser Asp
50 55 60

Leu Asp Tyr Leu Lys Ile Ser Phe Thr Glu Lys Gly Lys Gln Glu Asn
65 70 75 80

Ile Gln Lys Trp Ile Asn Gly Met Ser Trp Gly Ile Val Phe Tyr Lys
85 90 95

Tyr Gly Gly Gly Ala Gly Ser Thr Leu Thr Ile Arg Leu Arg Ile Glu
100 105 110

Thr Gly Thr Glu Pro Pro Val Ala Val Gly Pro Asp Lys Val Leu Ala
115 120 125

Glu Gln Gly Pro Pro Ala Leu Glu Pro Pro His Asn Leu Pro Val Pro
130 135 140

Gln Leu Thr Ser Leu Arg Pro Asp Ile Thr Gln Pro Pro Ser Asn Gly
145 150 155 160

Thr Thr Gly Leu Ile Pro Thr Asn Thr Pro Arg Asn Ser Pro Gly Val
165 170 175

Pro Val Lys Thr Gly Gln Arg Leu Phe Ser Leu Ile Gln Gly Ala Phe
180 185 190

Gln Ala Ile Asn Ser Thr Asp Pro Asp Ala Thr Ser Ser Cys Trp Leu
195 200 205

Cys Leu Ser Ser Gly Pro Pro Tyr Tyr Glu Gly Met Ala
210 215 220

<210> 11

<211> 220

<212> PRT

<213> porcine endogenous retrovirus Type PoEV1

<400> 11

Glu Lys Glu Lys Tyr Cys Gly Gly Ser Gly Glu Ser Phe Cys Arg Arg
 1 5 10

Trp Ser Cys Val Thr Ser Asn Asp Gly Asp Trp Lys Trp Pro Ile Ser
 20 25 30

Leu Gln Asp Arg Val Lys Phe Ser Phe Val Asn Ser Gly Pro Gly Lys
 35 40 45

Tyr Lys Met Met Lys Leu Tyr Lys Asp Lys Ser Cys Ser Pro Ser Asp
 50 55 60

Leu Asp Tyr Leu Lys Ile Ser Phe Thr Glu Arg Lys Thr Gly Lys Tyr
 65 70 75 80

Ser Lys Val Asp Lys Trp Tyr Glu Leu Gly Asn Ser Phe Leu Leu Tyr
 85 90 95

Gly Gly Gly Ala Gly Ser Thr Leu Thr Ile Arg Leu Arg Ile Glu Thr
 100 105 110

Gly Thr Glu Pro Pro Val Ala Met Gly Pro Asp Lys Val Leu Ala Glu
 115 120 125

Gln Gly Pro Pro Ala Leu Glu Pro Pro His Asn Leu Pro Val Pro Gln
 130 135 140

Leu Thr Ser Leu Arg Pro Asp Ile Thr Gln Pro Pro Ser Asn Ser Thr
 145 150 155 160

Thr Gly Leu Ile Pro Thr Asn Thr Pro Arg Asn Ser Pro Gly Val Pro
 165 170 175

Val Lys Thr Gly Gln Arg Leu Phe Ser Leu Ile Gln Gly Ala Phe Gln
 180 185 190

Ala Ile Asn Ser Thr Asp Pro Asp Ala Thr Ser Ser Cys Trp Leu Cys
 195 200 205

Leu Ser Ser Gly Pro Pro Tyr Tyr Glu Gly Met Ala
 210 215 220

<210> 12

<211> 15

<212> PRT

<213> porcine endogenous retrovirus

<400> 12

Arg Glu Glu Arg Arg Asp Arg Arg Gln Glu Lys Asn Leu Thr Lys
1 5 10 15

<210> 13

<211> 14

<212> PRT

<213> porcine endogenous retrovirus

<400> 13

Ala Arg Asn Cys Pro Lys Lys Gly Asn Lys Gly Pro Lys Val
1 5 10

<210> 14

<211> 16

<212> PRT

<213> porcine endogenous retrovirus Type PoEV1

<400> 14

Thr Ser Leu Arg Pro Asp Ile Thr Gln Pro Pro Ser Asn Ser Thr Thr
1 5 10 15

<210> 15

<211> 15

<212> PRT

<213> porcine endogenous retrovirus Type PERV B

<400> 15

Lys Gly Lys Gln Glu Asn Ile Gln Lys Trp Ile Asn Gly Met Ser
1 5 10 15

<210> 16

<211> 18

<212> PRT

<213> porcine endogenous retrovirus Type PoEV1

<400> 16

Arg Lys Thr Gly Lys Tyr Ser Lys Val Asp Lys Trp Tyr Glu Leu Gly
1 5 10 15

Asn Ser

<210> 17

<211> 12

<212> PRT

<213> porcine endogenous retrovirus Type Tsukuba

<400> 17

Asn Thr Val Leu Thr Gly Gln Arg Pro Pro Thr Gln
1 5 10

<210> 18

<211> 22

<212> PRT

<213> porcine endogenous retrovirus Type PERV A

<400> 18

Gly His Gly Arg Trp Lys Asp Trp Gln Gln Arg Val Gln Lys Asp Val
1 5 10 15

Arg Asn Lys Gln Ile Ser
20

<210> 19

<211> 16

<212> PRT

<213> porcine endogenous retrovirus Type PERV A

<400> 19

Ile Gln Glu Gln Arg Pro Ser Pro Asn Pro Ser Asp Tyr Asn Thr Thr
1 5 10 15